arXiv:math/0304140v5 [math.AG] 21 Aug 2003

TRANSACTIONS OF THE
AMERICAN MATHEMATICAL SOCIETY
Volume 00, Number 0, Pages 000–000
S 0002-9947(XX)0000-0

THE CHEN-RUAN COHOMOLOGY OF WEIGHTED PROJECTIVE SPACES

YUNFENG JIANG

Abstract. Chen and Ruan [6] defined a very interesting cohomology theory for orbifolds, which is now called Chen-Ruan cohomology. The primary objective of this paper is to compute the Chen-Ruan cohomology rings of the weighted projective spaces, a class of important spaces in physics. The classical tools (Chen-Ruan cohomology, toric varieties, the localization technique) which have been proved to be successful are used to study the orbifold cohomology of weighted projective spaces. Given a weighted projective space $\mathbf{P}^n_{q_0,\cdots,q_n}$, we determine all of its twisted sectors and the corresponding degree shifting numbers, and we calculate the orbifold cohomology group of $\mathbf{P}^n_{q_0,\cdots,q_n}$. For a general reduced weighted projective space, we give a formula to compute the 3-point function which is the key in the definition of Chen-Ruan cohomology ring. Finally we concretely calculate the Chen-Ruan cohomology ring of weighted projective space $\mathbf{P}^5_{1,2,2,3,3,3}$.

1. Introduction

The notion of Chen-Ruan orbifold cohomology has appeared in physics as a result of studying the string theory on global quotient orbifold, (see [9] and [10]). In addition to the usual cohomology of the global quotient, this space included the cohomology of so-called twisted sectors. Zaslow [25] gave a lot of examples of global quotients and computed their orbifold cohomology spaces. But the real mathematical definition of orbifold cohomology was given by Chen and Ruan [6] for arbitrary orbifolds. The most interesting feature of this new cohomology theory, besides the generalization of non global quotients, is the existence of a ring structure which was previously missing. This ring structure is obtained from Chen-Ruan's orbifold quantum cohomology construction (see [7]) by restricting to the class called ghost maps, the same as the ordinary cup product may be obtained by quantum cup product. Since the Chen-Ruan cohomology appeared, the problem of how to calculate the orbifold cohomology has been considered by several authors. Chen and Ruan [6] gave several simple examples. Chen Hao [4] computed the orbifold cohomology group of moduli space $\mathcal{M}_{0,n}/S_n$. B.Doug Park and Mainak Poddar [22] considered the Chen-Ruan cohomology ring of the mirror quintic. All the above examples are orbifold global quotients. In this paper we calculate the Chen-Ruan cohomology rings of weighted projective spaces-a large class of non-global quotient orbifolds.

Key words and phrases. Chen-Ruan cohomology, twisted sectors, toric varieties, weighted projective space, localization.

©1997 American Mathematical Society

A very power tool to compute the Chen-Ruan cohomology of weighted projective spaces is the method of toric varieties. The theory of toric varieties establishes a classical connection between algebraic geometry and the theory of convex polytopes. From the fan of a toric variety, we can obtain a lot of information about the toric variety. In particular, when the fan Σ of a toric variety X is simplicial, the toric variety X is an orbifold with finite abelian groups as local groups. In this paper, we take the weighted projective spaces as simplicial toric varieties with local isotropy groups the finite cyclic groups. And then using the properties of toric varieties induced from the fans, we calculate the Chen-Ruan cohomology group of any weighted projective space.

To calculate the Chen-Ruan cohomology ring of the weighted projective space, we use the Riemann bilinear relations for periods [15] to identify the obstruction bundle. Up to now, except that the obstruction bundle for the mirror quintic example calculated by Park and Poddar [22] is a nontrivial line bundle, all the other calculated obstruction bundles of the examples are trivial. In this paper, the obstruction bundle we consider is the Whitney sum of some line bundles, generalizing the case of mirror quintic example. And we also introduce the localization techniques [3] which should work for toric varieties to compute the 3-point function which is the key in the orbifold cup product [6]. In particular, we give a concrete example.

On the other hand, a very interesting aspect of calculating the Chen-Ruan cohomology rings of weighted projective spaces lies in a conjecture of Ruan. In string theory, physicists suggest that the orbifold string theory of an orbifold should be equivalent to the ordinary string theory of its crepant resolution. For the orbifold cohomology, the Cohomology Hyperkahler Resolution Conjecture of Ruan (see [23]) states that the Chen-Ruan cohomology ring of an orbifold should be isomorphic to the ordinary cohomology ring of its hyperkahler resolution. I hope that my calculation of the Chen-Ruan cohomology ring of the weighted projective space may contribute to this interesting problem.

The thesis is outlined as follows. Section 2 is a review of some basic facts concerning orbifold, Chen-Ruan cohomology and simplicial toric varieties. In section 3 we introduced the basic concept of the weighted projective space. In section 4 we discuss the Chen-Ruan cohomology group of any weighted projective space. And in the section 5 we compute the ring structure of the Chen-Ruan cohomology of the weighted projective space.

2. Preliminaries

2.1. **Orbifold and orbifold vector bundle.**

Definition 2.1.1. An orbifold structure on a Hausdorff, separate topological space X is given by an open cover $\mathcal{U}$ of X satisfying the following conditions.

(1) Each element U in $\mathcal{U}$ is uniformized, say by (V, G, π). Namely, V is a smooth manifold and G is a finite group acting smoothly on V such that $U = V/G$ with π as the quotient map. Let $Ker(G)$ be the subgroup of G acting trivially on V.

(2) For $U^{'} \subset U$, there is a collection of injections $(V^{'}, G^{'}, \pi^{'}) \longrightarrow (V, G, \pi)$. Namely, the inclusion $i : U^{'} \subset U$ can be lifted to maps $\widetilde{i} : V^{'} \longrightarrow V$ and an injective homomorphism $i_{*} : G^{'} \longrightarrow G$ such that i_{*} is an isomorphism from $Ker(G^{'})$ to $Ker(G)$ and $\widetilde{i}$ is i_{*}-equivariant.

(3) For any point $x \in U_1 \cap U_2$, $U_1, U_2 \in \mathcal{U}$, there is a $U_3 \in \mathcal{U}$ such that $x \in U_3 \subset U_1 \cap U_2$.

For any point $x \in X$, suppose that (V, G, π) is a uniformizing neighborhood and $\overline{x} \in \pi^{-1}(x)$. Let G_x be the stabilizer of G at $\overline{x}$. Up to conjugation, it is independent of the choice of $\overline{x}$ and is called the *local group* of x. Then there exists a sufficiently small neighborhood V_x of $\overline{x}$ such that (V_x, G_x, π_x) uniformizes a small neighborhood of x, where π_x is the restriction $\pi \mid V_x$. (V_x, G_x, π_x) is called *a local chart* at x. The orbifold structure is called *reduced* if the action of G_x is effective for every x.

Let $pr : E \longrightarrow X$ be a rank k complex *orbifold bundle* over an orbifold X([6]). Then a uniformizing system for $E \mid U = pr^{-1}(U)$ over a uniformized subset U of X consists of the following data:

(1) A uniformizing system (V, G, π) of U.

(2) A uniformizing system $(V \times \mathbf{C^k}, G, \widetilde{\pi})$ for $E \mid U$. The action of G on $V \times \mathbf{C}^k$ is an extension of the action of G on V given by $g \cdot (x, v) = (g \cdot x, \rho(x, g)v)$ where $\rho : V \times G \longrightarrow Aut(\mathbf{C}^k)$ is a smooth map satisfying:

$$\rho(g \cdot x, h) \circ \rho(x, g) = \rho(x, hg), g, h \in G, x \in V.$$

(3)The natural projection map $\widetilde{pr} : V \times \mathbf{C^k} \longrightarrow \mathbf{V}$ satisfies $\pi \circ \widetilde{pr} = pr \circ \widetilde{\pi}$.

By an orbifold connection $\triangle$ on E we mean an equivariant connection that satisfies $\triangle = g^{-1} \triangle g$ for every uniformizing system of E. Such a connection can be always obtained by averaging an equivariant partition of unity.

2.2. **Twisted Sectors and Chen-Ruan Cohomology.** The most physical idea is twisted sectors. Let X be an orbifold. Consider the set of pairs:

$$\widetilde{X}_k = \{(p, (\mathbf{g})_{G_p}) | p \in X, \mathbf{g} = (g_1, \cdots, g_k), g_i \in G_p\}$$

where $(\mathbf{g})_{G_p}$ is the conjugacy class of k-tuple $\mathbf{g} = (g_1, \cdots, g_k)$ in G_p. We use G^k to denote the set of k-tuples. If there is no confusion, we will omit the subscript G_p to simplify the notation. Suppose that X has an orbifold structure $\mathcal{U}$ with uniformizing systems $(\widetilde{U}, G_U, \pi_U)$. From Chen and Ruan [6], also see [18], we have: $\widetilde{X}_k$ is naturally an orbifold, with the generalized orbifold structure at $(p, (\mathbf{g})_{G_p})$ given by $(V_p^{\mathbf{g}}, C(\mathbf{g}), \pi : V_p^{\mathbf{g}} \longrightarrow V_p^{\mathbf{g}}/C(\mathbf{g}))$, where $V_p^{\mathbf{g}} = V_p^{g_1} \cap \cdots V_p^{g_k}$, $C(\mathbf{g}) = C(g_1) \cap \cdots C(g_k)$. Here $\mathbf{g} = (g_1, \cdots, g_k)$, V_p^g stands for the fixed point set of g in V_p. When X is almost complex, $\widetilde{X_k}$ inherits an almost complex structure from X, and when X is closed, $\widetilde{X_k}$ is finite disjoint union of closed orbifolds.

Now we describe the the connected components of $\widetilde{X_k}$, Recall that every point p has a local chart (V_p, G_p, π_p) which gives a local uniformized neighborhood $U_p = \pi_p(V_p)$. If $q \in U_p$, up to conjugation there is a unique injective homomorphism $i_* : G_q \longrightarrow G_p$. For $\mathbf{g} \in (G_q)^k$, the conjugation class $i_*(\mathbf{g})_q$ is well defined. We define an equivalence relation $i_*(\mathbf{g})_q \cong (\mathbf{g})_q$. Let T_k denote the set of equivalence classes.To abuse the notation, we use $(\mathbf{g})$ to denote the equivalence class which $(\mathbf{g})_q$ belongs to. We will usually denote an element of T_1 by (g). It is clear that $\widetilde{X_k}$ can be decomposed as a disjoint union of connected components:

$$\widetilde{X_k} = \bigsqcup_{(\mathbf{g}) \in T_k} X_{(\mathbf{g})}$$

Where $X_{(\mathbf{g})} = \{(p, (\mathbf{g}^{'})_p) | \mathbf{g}^{'} \in (G_p)^k, (\mathbf{g}^{'})_p \in (\mathbf{g})\}$. Note that for $\mathbf{g} = (1, \cdots, 1)$, we have $X_{(\mathbf{g})} = X$. A component $X_{(\mathbf{g})}$ is called a $k - multisector$, if $\mathbf{g}$ is not the identity. A component of $X_{(g)}$ is simply called a *twisted sector*. If X has an almost complex, complex or kahler structure, then $X_{(\mathbf{g})}$ has the analogous structure induced from X. We define

$$T_3^0 = \{(\mathbf{g}) = (g_1, g_2, g_3) \in T_3 | g_1 g_2 g_3 = 1\}.$$

Note that there is an one to one correspondence between T_2 and T_3^0 given by $(g_1, g_2) \longmapsto (g_1, g_2, (g_1 g_2)^{-1})$.

Now we define the Chen-Ruan cohomology. Assume that X is a n-dimensional compact almost complex orbifold with almost structure J. Then for a point p with nontrivial group G_p, J gives rise to an effective representation $\rho_p : G_p \longrightarrow GL(n, \mathbf{C})$. For any $g \in G_p$, we write $\rho_p(g)$, up to conjugation, as a diagonal matrix

$$diag\left(e^{2\pi i \frac{m_{1,g}}{m_g}}, \ldots, e^{2\pi i \frac{m_{n,g}}{m_g}}\right).$$

where m_g is the order of g in G_p, and $0 \leq m_{i,g} < m_g$. Define a function $\iota : \widetilde{X_1} \longrightarrow \mathbf{Q}$ by

$$\iota(p, (g)_p) = \sum_{i=1}^{n} \frac{m_{i,g}}{m_g}.$$

We can see that the function $\iota : \widetilde{X_1} \longrightarrow \mathbf{Q}$ is locally constant and $\iota = 0$ if $g = 1$. Denote its value on $X_{(g)}$ by ι_g. We call ι_g the degree shifting number of $X_{(g)}$. It has the following properties:

(1) $\iota_{(g)}$ is an integer iff $\rho_p(g) \in SL(n, \mathbf{C})$;

(2) $\iota_{(g)} + \iota_{(g^{-1})} = rank(\rho_p(g) - Id) = n - dim_{\mathbf{C}} X_{(g)}$.

A C^∞ differential form on X is a G-invariant differential form on V for each uniformizing system (V, G, π). Then orbifold integration is defined as follows. Suppose $U = V/G$ is connected, for any compactly supported differential n-form ω on U, which is, by definition, a G-invariant n-form $\widetilde{\omega}$ on V,

$$\int_U^{orb} \omega := \frac{1}{|G|} \int_V \widetilde{\omega} \tag{2.1}$$

Where $|G|$ is the order of G. The orbifold integration over X is defined by using a C^∞ partition of unity. The orbifold integration coincides with the usual measure theoretic integration iff the orbifold structure is reduced.

Holomorphic forms for a complex orbifold X are again obtained by patching G-invariant holomorphic forms on the uniformizing system (V, G, π). We consider the Cech cohomology groups of X and $X_{(\mathbf{g})}$ with coefficients in the sheaves of holomorphic forms. The Cech cohomology groups can be identified with the Dolbeault cohomology groups of (p, q)-forms [2].

Definition 2.2.1. ([6]) Let X be a closed complex orbifold, we define the orbifold cohomology group of X by

$$H^d_{orb} := \bigoplus_{(g) \in T_1} H^{d - 2\iota_{(g)}}(X_{(g)}, \mathbf{Q})$$

For $0 \leq p, q \leq dim_{\mathbf{C}} X$, we define the orbifold Dolbeault cohomology group of X by

$$H^{p,q}_{orb}(X) := \bigoplus_{(g)\in T_1} H^{p-\iota_{(g)},q-\iota_{(g)}}(X_{(g)},\mathbf{C})$$

2.3. **The Obstruction Bundle.** Choose $(\mathbf{g})= (g_1, g_2, g_3) \in T_3^0$. Let $(p, (\mathbf{g})_p)$ be a generic point in $X_{(\mathbf{g})}$. Let $K(\mathbf{g})$ be the subgroup of G_p generated by g_1 and g_2. Consider an orbifold Riemann sphere with three orbifold points $(S^2, (p_1, p_2, p_3), (k_1, k_2, k_3))$. When there is no confusion, we will simply denote it by S^2. The orbifold fundamental group is:

$$\pi_1^{orb}(S^2) = \{\lambda_1, \lambda_2, \lambda_3 | \lambda_i^{k_i} = 1, \lambda_1\lambda_2\lambda_3 = 1\}$$

Where λ_i is represented by a loop around the marked p_i. There is a surjective homomorphism

$$\rho : \pi_1^{orb}(S^2) \longrightarrow K(\mathbf{g})$$

specified by mapping $\lambda_i \longmapsto g_i$. $Ker(\rho)$ is a finite-index subgroup of $\pi_1^{orb}(S^2)$. Let $\widetilde{\Sigma}$ be the orbifold universal cover of S^2. Let $\Sigma = \widetilde{\Sigma}/Ker(\rho)$. Then Σ is smooth, compact and $\Sigma/K(\mathbf{g}) = S^2$. The genus of Σ can be computed using Riemann Hurwitz formula for Euler characteristics of a branched covering, and turns out to be

$$g(\Sigma) = \frac{1}{2}(2 + |K(\mathbf{g})| - \Sigma_{i=1}^3 \frac{|K(\mathbf{g})|}{k_i}) \tag{2.2}$$

$K(\mathbf{g})$ acts holomorphically on Σ and hence $K(\mathbf{g})$ acts on $H^{0,1}(\Sigma)$. The "obstruction bundle" $E_{(\mathbf{g})}$ over $X_{(\mathbf{g})}$ is constructed as follows. On the local chart $(V_p^{\mathbf{g}}, C(\mathbf{g}), \pi)$ of $X_{(\mathbf{g})}$, $E_{(\mathbf{g})}$ is given by $(TV_p \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})} \times V_p^{\mathbf{g}} \longrightarrow V_p^{\mathbf{g}}$, where $(TV_p \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})}$ is the $K(\mathbf{g})$-invariant subspace. We define an action of $C(\mathbf{g})$ on $TV_p \otimes H^{0,1}(\Sigma)$, which is the usual one on TV_p and trivial on $H^{0,1}(\Sigma)$. The the action of $C(\mathbf{g})$ and $K(\mathbf{g})$ commute and $(TV_p \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})}$ is invariant under $C(\mathbf{g})$. Thus we have obtained an action of $C(\mathbf{g})$ on $(TV_p \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})} \times V_p^{\mathbf{g}} \longrightarrow V_p^{\mathbf{g}}$, extending the usual one on $V_p^{\mathbf{g}}$. These trivializations fit together to define the bundle $E_{(\mathbf{g})}$ over $X_{(\mathbf{g})}$. If we set $e : X_{(\mathbf{g})} \longrightarrow X$ to be the map given by $(p, (\mathbf{g})_p) \longmapsto p$, one may think of $E_{(\mathbf{g})}$ as $(e^*TX \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})}$. The rank of $E_{(\mathbf{g})}$ is given by the formula [6]:

$$rank_{\mathbf{C}}(E_{(\mathbf{g})}) = dim_{\mathbf{C}}(X_{(\mathbf{g})}) - dim_{\mathbf{C}}(X) + \Sigma_{j=1}^3 \iota_{(g_j)} \tag{2.3}$$

2.4. **Orbifold cup product.** First, there is a natural map $I : X_{(g)} \longrightarrow X_{(g^{-1})}$ defined by $(p, (g)_p) \longmapsto (p, (g^{-1})_p)$.

Definition 2.4.1. Let $n = dim_{\mathbf{C}}(X)$. For any integer $0 \le n \le 2n$, the pairing

$$<,>_{orb}: H^d_{orb}(X) \times H^{2n-d}_{orb}(X) \longrightarrow \mathbf{Q}$$

is defined by taking the direct sum of

$$<,>^{(g)}_{orb}: H^{d-2\iota_{(g)}}(X_{(g)}; \mathbf{Q}) \times H^{2n-d-2\iota_{(g^{-1})}}(X_{(g^{-1})}; \mathbf{Q}) \longrightarrow \mathbf{Q}$$

where

$$< \alpha, \beta >^{(g)}_{orb} = \int^{orb}_{X_{(g)}} \alpha \wedge I^*(\beta)$$

for $\alpha \in H^{d-2\iota_{(g)}}(X_{(g)};\mathbf{Q})$, and $\beta \in H^{2n-d-2\iota_{(g^{-1})}}(X_{(g^{-1})};\mathbf{Q})$.

Choose an orbifold connection A on $E_{(\mathbf{g})}$. Let $e_A(E_{(\mathbf{g})})$ be the Euler form computed from the connection A by Chen-Weil theory. Let $\eta_j \in H^{d_j}(X_{(g_j)};\mathbf{Q})$, for $j = 1,2,3$. Define maps $e_j : X_{(\mathbf{g})} \longrightarrow X_{(g_j)}$ by $(p,(\mathbf{g})_p) \longmapsto (p,(g_j)_p)$.

Definition 2.4.2. Define the 3-point function to be

$$< \eta_1, \eta_2, \eta_3 >_{orb} := \int^{orb}_{X_{(\mathbf{g})}} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge e_A(E_{(\mathbf{g})}) \tag{2.4}$$

Note that the above integral does not depend on the choice of A. As in the definition 2.4.1, we extend the 3-point function to $H^*_{orb}(X)$ by linearity. We define the orbifold cup product by the relation

$$< \eta_1 \cup_{orb} \eta_2, \eta_3 >_{orb} := < \eta_1, \eta_2, \eta_3 >_{orb} \tag{2.5}$$

Again we extend $\cup_{orb}$ to $H^*_{orb}(X)$ via linearity. Note that if $(\mathbf{g}) = (1,1,1)$, then $\eta_1 \cup_{orb} \eta_2$ is just the ordinary cup product $\eta_1 \cup \eta_2$ in $H^*(X)$.

2.5. **Simplicial Toric Varieties as Orbifolds.**

A toric variety is a normal variety with an action of an algebraic torus which admits an open dense orbit homeomorphic to the torus. Every toric variety is described by a set of combinatoric data, called a fan Ξ in a lattice N, [13], [20]. Ξ is simplicial if every cone σ in Ξ is generated by a subset of a basis of $\mathbf{R}^n = N \otimes \mathbf{R}$.

We now describe the orbifold structure of simplicial toric varieties, see Poddar [20]. Let Ξ be any simplicial fan in a n dimensional lattice N. X_Ξ be the corresponding toric variety. For a cone $\tau \in \Xi$, denote the set of its primitive 1-dimensional generators by $\tau[1]$, the corresponding affine open subset of X_Ξ by U_τ, and the corresponding torus orbit by O_τ. We write $\nu \leq \tau$ if the cone ν is a face of the cone τ, and $\nu < \tau$ if it is a proper subface. $U_\tau = \sqcup_{\nu \leq \tau} O_\nu$. Let $M = Hom(N,\mathbf{Z})$ be the dual lattice of N with dual pair $<,>$. For any cone $\tau \in \Xi$, denote its dual cone in $M \otimes \mathbf{R}$ by $\check{\tau}$. Let $S_\tau = \check{\tau} \cap M$. $\mathbf{C}(\mathbf{S}_\tau)$ is the $\mathbf{C}$-algebra with generators χ^m for each $m \in S_\tau$ and relation $\chi^m \chi^{m'} = \chi^{m+m'}$. $U_\tau = Spec(\mathbf{C}[S_\tau])$.

Then the orbifold structure of the toric variety X_Ξ can be described as follows. Let σ be any n dimensional cone of Ξ. Let $v_1, \cdots, v_n$ be the primitive 1 dimension generstors of σ. These are linearly independent in $N_\mathbf{R} = N \otimes \mathbf{R}$. Let N_σ be the sublattice of N generated by $v_1, \cdots, v_n$. And let $G_\sigma = N/N_\sigma$ be the quotient group, then G_σ is finite and abelian.

Let σ' be the cone σ regarded in N_σ. Let $\check{\sigma}'$ be the dual cone of σ' in M_σ, the dual lattice of N_σ. $U_{\sigma'} = spec(\mathbf{C}[\check{\sigma}' \cap M_\sigma])$. Note that σ' is a smooth cone in N_σ. So $U_{\sigma'} \cong \mathbf{C}^n$.

Now there is a canonical dual pairing $M_\sigma/M \times N/N_\sigma \longrightarrow \mathbf{Q}/\mathbf{Z} \longrightarrow \mathbf{C}^*$, the first map by the pairing $<,>$ and the second by $q \longmapsto exp(2\pi i q)$. Now G_σ acts on $\mathbf{C}[M_\sigma]$, the group ring of M_σ, by: $v(\chi^u) = exp(2\pi i < u, v >)\chi^u$, for $v \in N$ and $u \in M_\sigma$. Note that

$$(\mathbf{C}[M_\sigma])^{G_\sigma} = \mathbf{C}[M] \tag{2.6}$$

Thus G_σ acts on $U_{\sigma'}$. Let π_σ be the quotient map. Then $U_\sigma = U_{\sigma'}/G_\sigma$. So U_σ is uniformized by $(U_{\sigma'}, G_\sigma, \pi_\sigma)$. For any $\tau < \sigma$, the orbifold structure on U_τ is the same as the one induced from the uniformizing system on U_σ. Then by the description of the toric gluing it is clear that $\{(U_{\sigma'}, G_\sigma, \pi_\sigma) : \sigma \in \Xi[n]\}$ defines a reduced orbifold structure on X_Ξ. We give a more explicit verification of this fact below.

Let B be the nonsingular matrix with generators $v_1, \cdots, v_n$ of σ as rows. Then $\check{\sigma}'$ is generated in M_σ by the column vectors $v^1, \cdots, v^n$ of the matrix B^{-1}. So $\chi^{v^1}, \cdots, \chi^{v^n}$ are the coordinates of $U_{\sigma'}$. For any $k = (k_1, \cdots, k_n) \in N$, the corresponding coset $[k] \in G_\sigma$ acts on $U_{\sigma'}$ in these coordinates as a diagonal matrix: $diag(exp(2\pi i c_1), \cdots, exp(2\pi i c_n))$, where $c_i =< k, v^i >$. Such a matrix is uniquely represented by an n-tuple $a = (a_1, \cdots, a_n)$ where $a_i \in [0,1)$ and $c_i = a_i + b_i, b_i \in \mathbf{Z}$. In matrix notation, $kB^{-1} = a + b \cong k = aB + bB$. We denote the integral vector aB in N by k_a and the diagonal matrix corresponding to a by g_a. $k_a \longleftrightarrow g_a$ gives a one to one correspondence between the elements of G_σ and the integral vector in N that are linear combinations of the generators of σ with coefficient in $[0,1)$.

Now let us examine the orbifold chart induced by $(U_{\sigma'}, G_\sigma, \pi_\sigma)$ at any point $p \in U_\sigma$. By the orbit decomposition, there is a unique $\tau \in \Xi$ such that $p \in O_\tau$. We assume τ is generated by $v_1, \cdots, v_j, j \leq n$. Then any preimage of p with respect to π_σ has coordinates $\chi^{v^i} = 0$ iff $i \leq j$. Let $z = (0, \ldots, 0, z_{j+1}, \ldots, z_n)$ be one such preimage. Let $G_\tau := \{g_a \in G_\sigma : a_i = 0 \ if \ j+1 \leq i \leq n\}$. We can find a small neighborhood $W \subset (\mathbf{C}^*)^{n-j}$ of $(z_{j+1}, \ldots, z_n)$ such that the inclusions $\mathbf{C}^j \times W \hookrightarrow U_{\sigma'}$ and $G_\tau \hookrightarrow G_\sigma$ induces an injection of uniformizing systems $(\mathbf{C}^j \times W, G_\tau, \pi) \hookrightarrow (U_{\sigma'}, G_\sigma, \pi_\sigma)$ on some small open neighborhood U_p of p. So we have $G_p = G_\tau$ and an orbifold chart $(\mathbf{C}^j \times W, G_\tau, \pi)$. Note that G_τ can be constructed from the set $\{k_a = \Sigma_{i=1}^{j} a_i v_i : k_a \in N, a_i \in [0,1)\}$ which is completely determined by τ and hence is independent of σ.

3. The Weighted Projective Spaces

3.1. **The Definition and the Orbifold Structure of the Weighted Projective Space.** Throughout this paper, $a(\mathbf{p})$ for $a \in \mathbf{C}$ and $\mathbf{p} = (p_1, \cdots, p_n) \in \mathbf{Z}^n$ will denote the diagonal matrix:

$$Diag(a^{p_1}, \cdots, a^{p_n})$$

with diagonal entries $a^{p_i}, i = 1, \cdots, n$. Moreover, for an integer q, μ_q will denote the group $\mathbf{Z}/q\mathbf{Z}$.

Definition 3.1.1. ([14]) Let $Q = (q_0, \cdots, q_n)$ be a $(n+1)$-tuple of positive integers. The weighted projective space of type Q, $\mathbf{P}^n(Q) = \mathbf{P}^n_{q_0, \cdots, q_n}$ is defined by

$$\mathbf{P}^n_{q_0, \cdots, q_n} = \left\{ z \in (\mathbf{C}^{n+1})^* | z \sim \lambda(\mathbf{q}) \cdot z, \lambda \in \mathbf{C}^* \right\}$$

where $\lambda(\mathbf{q}) = Diag(\lambda^{q_0}, \cdots, \lambda^{q_n})$.

Remark 3.1.2. (1)The above $\mathbf{C}^*$-action is free iff $q_i = 1$ for every $i = 0, \cdots, n$; (2) If $gcd(q_0, \cdots, q_n) = d \neq 1$, then $\mathbf{P}^n_{q_0, \cdots, q_n}$ is homeomorphic to $\mathbf{P}^n_{q_0/d, \cdots, q_n/d}$(by identification of λ^d with λ).

Weighted projective spaces are, in general, orbifolds where the singularities have cyclic structure groups acting diagonally. Moreover, if all the $q_i' s$ are mutually prime, all these orbifold singularities are isolated. In fact, as is usually done for complex projective spaces, we can consider the sets

$$U_i = \left\{ [z]_Q \in \mathbf{P}^n_{q_0, \cdots, q_n} : z_i \neq 0 \right\} \subset \mathbf{P}^n_{q_0, \cdots, q_n}$$

and the bijective maps ϕ_i from U_i to $\mathbf{C}^n/\mu_{q_i}(Q_i)$ given by

$$\phi_i([z]_Q) = \left(\frac{z_0}{(z_i)^{q_0/q_i}}, \cdots, \frac{\hat{z_i}}{z_i}, \cdots, \frac{z_n}{(z_i)^{q_n/q_i}} \right)_{q_i}$$

where $(z_i)^{1/q_i}$ is a q_i-root of z_i and $(\cdot)_{q_i}$ is a μ_i-conjugacy class in $\mathbf{C}^n/\mu_{q_i}(Q_i)$ with μ_{q_i} acting on $\mathbf{C}^n$ by

$$\xi \cdot z = \xi(Q_i)z, \xi \in \mu_{q_i}$$

Here $Q_i = (q_0, \cdots, \hat{q_i}, \cdots, q_n)$. Then on $\phi_i(U_j \cap U_i) \subset \mathbf{C}^n/\mu_{q_i}(Q_i)$,

$$\phi_j \circ \phi_i^{-1}((z_1, \cdots, z_n)_{q_i}) = \left(\frac{z_0}{(z_j)^{q_0/q_j}}, \cdots, \frac{\hat{z_j}}{z_j}, \cdots, \frac{1}{(z_j)^{q_i/q_j}}, \cdots, \frac{z_n}{(z_j)^{q_n/q_j}} \right)_{q_j}$$

3.2. **Toric Structure of the Weighted Projective Spaces.** Given $Q = (q_0, \cdots, q_n) \in \mathbf{Z}^{n+1}$, let $d_{i_1,\cdots,i_s} = gcd(q_{i_1}, \cdots, q_{i_s})$ and $d_j = gcd(q_0, \cdots, \hat{q}_j, \cdots, q_n)$ $(i_1, \cdots, i_s, j \in \{1, \cdots, n\})$. Define a grading of $\mathbf{C}[X_0, \cdots, X_n]$ by $degX_i = q_i$. We denote this ring by $S(Q)$. Then $\mathbf{P}^n_{q_0,\cdots,q_n} = projS(Q)$ is the weighted projective space of type Q. $\mathbf{P}^n_{q_0,\cdots,q_n}$ is covered by the affine open sets $D_+(X_i) := specS(Q)_{X_i}, (i = 0, ..., n)$. The monic monomials of $S(Q)_{X_i}$ are of type $X_i^{-l}\Pi_{j\neq i}X_j^{\lambda_j}$, where $lq_i = \Sigma_{j\neq i}\lambda_j q_j$ and l, λ_j are non-negative integers. So each such monomial is uniquely determined by the n-tuple $(\lambda_0, \cdots, \lambda_{i-1}, \lambda_{i+1}, \cdots, \lambda_n)$ of its non-negative exponents. The exponents occurring are just the points lying in the intersection of the cone $e := pos\{e_1, \cdots, e_n\}$ and the lattice $N_{Q,q_i} \subseteq \mathbf{Z}^n$ that is defined as follows.

Consider $Q_i = (q_0, \cdots, \hat{q_i}, \cdots, q_n)$ as an element of $Hom_{\mathbf{Z}}(\mathbf{Z}^n, \mathbf{Z})$ by setting:

$$Q_i(a_1, \cdots, a_n) := q_0 a_1 + \cdots, +q_n a_n$$

$\mathbf{Z}^n$ being equipped with its canonical basis. Let $\pi_i : \mathbf{Z} \longrightarrow \mathbf{Z}_{q_i}$ denote the canonical projection. Then

$$N_{Q,q_i} := Ker(\mathbf{Z}^n \longrightarrow \mathbf{Z} \longrightarrow \mathbf{Z}_{q_i})$$

is a sublattice of $\mathbf{Z}^n$. Denote by $M_{Q,i}$ the dual lattice. We have an isomorphism of semigroup rings

$$S(Q)_{(X_i)} \cong \mathbf{C}[e \cap N_{Q,q_i}]$$

revealing $D_+(X_i)$ to be the affine toric variety associated with $\check{e}$ with respect to M_{Q,q_i}.

Proposition 3.2.1. *([5]) Let $C_i = (c^i_1, \cdots, c^i_n)$ be a basis of N_{Q,q_i} and denote by $r^i_1, \cdots, r^i_n$ the row vectors of C_i. Let $\sigma_i := pos\{r^i_1, \cdots, r^i_n\}$. Then there is an isomorphism of semigroups*

$$\check{\sigma}_i \cap \mathbf{Z}^n \simeq e \cap N_{Q,q_i}.$$

Proposition 3.2.2. *([5]) With the natation introduced above, the matrix*

$$C_0 = \begin{pmatrix} \frac{q_0}{d_{01}} & c^0_{12} & \cdots & c^0_{1n} \\ 0 & \frac{d_{01}}{d_{012}} & \cdots & c^0_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \frac{d_n}{d_{0\cdots n}} \end{pmatrix} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix} = (c^0_{ij})$$

is a basis of N_{Q,q_0}*, where* $c^0_{ij} \in \mathbf{Z}_{\geq \mathbf{0}}$ *are determined as follows. For fixed* $j \in \{2, ..., n\}$ *construct* c^0_{ij} *successively for* $i = j-1, ..., 1$ *by requiring* $c^0_{ij} \in \mathbf{Z}_{\geq \mathbf{0}}$ *to be minimal with the property that*

$$c^0_{ij} q_i + \sum_{\nu=i+1}^{j} c^0_{\nu j} q_\nu \in gcd(q_0, ..., q_{i-1})\mathbf{Z} = d_{0 \cdots i-1}\mathbf{Z}.$$

Proposition 3.2.3. *([5]) With the notations introduced above, let* $v_0 := -\sum_{i=1}^{n} \frac{q_i}{q_0} v_i$*,* $\rho_i = \mathbf{R}_{\geq 0} v_i$*. Then the complete fan* Ξ *determined by* $\Xi[1] = \{\rho_0, \cdots, \rho_n\}$ *is the fan of the weighted projective space* $\mathbf{P}^n_{q_0, \cdots, q_n}$*.*

Remark 3.2.4. (1) The weighted projective space $\mathbf{P}^n_{q_0,\cdots,q_n}$ can also be constructed as follows. Given a fan $\Xi = \{u_0, \cdots, u_n\}$ so that $q_0 u_0 + q_1 u_1 + \cdots + q_n u_n = 0$, then the toric variety X_Ξ is the weighted projective space $\mathbf{P}^n_{q_0,\cdots,q_n}$. From above we can see that $\{v_0, \cdots, v_n\}$ satisfies the condition $q_0 v_0 + q_1 v_1 + \cdots + q_n v_n = 0$. In fact, the proposition 3.2.2 and 3.2.3 gave a method to compute the fan of weighted projective space of type Q. (2) If $gcd(q_0, \cdots, q_n) = d \neq 1$, from the construction of the matrix C_0, we can see that $\mathbf{P}^n_{q_0,\cdots,q_n}$ and $\mathbf{P}^n_{q_0/d,\cdots,q_n/d}$ have the same fans, so they are homeomorphic.

3.3. **Homogeneous Coordinate Representations of Weighted Projective Spaces Induced from Toric Varieties.** In this section we use the theorem of David Cox (see [8]) to represent the weighted projective space $\mathbf{P}^n_{q_0,\cdots,q_n}$ as the geometric quotient $(\mathbf{C^{n+1}})^*/\mathbf{C}^*$. From the theorem of David Cox in [7], for a fan Ξ, the toric variety X_Ξ is a geometric quotient $(\mathbf{C})^r \backslash Z/G$ iff Ξ is simplicial, where r is the number of the 1-primitive generators,Z is a subvariety of $\mathbf{C}^r$, G is some subgroup of $(\mathbf{C}^*)^r$. To give this representation for the weighted projective space, we must compute the space Z and the group G. Let $\Xi = \{v_0, \cdots, v_n\}$ be the fan of weighted projective space $\mathbf{P}^n_{q_0,\cdots,q_n}$. Then there are $n+1$ 1-dimensional primitive generators, which give variables $x_0, \cdots, x_n$. Furthermore, the maximal cones of the fan are generated by the n-element subsets of $\{v_0, \cdots, v_n\}$. It follows from [9] that

$$Z = V(x_0, \cdots, x_n) = \{(0, \cdots, 0)\} \subset \mathbf{C}^{n+1}.$$

Now we describe the group G. From [9],

$$G = \left\{ (\mu_0, \cdots, \mu_n) \in (\mathbf{C}^*)^{n+1} | \Pi_{i=1}^{n+1} \mu_i^{<m, v_i>} = 1, \text{for all } m \in \mathbf{Z}^n \right\}$$

However it suffices to let m be the standard basis elements $e_1, \cdots, e_n$. Thus $(\mu_0, \cdots, \mu_n) \in G$ iff

$$\Pi_{i=0}^{n} \mu_i^{<e_1, r_i>} = \Pi_{i=0}^{n} \mu_i^{<e_2, r_i>} = \cdots = \Pi_{i=0}^{n} \mu_i^{<e_n, r_i>} = 1 \qquad (3.1)$$

From proposition 3.2.3, we have the vectors $v_1, \cdots, v_n$, and $v_0 = -\Sigma_{i=1}^{n} \frac{q_i}{q_0} r_i$, so we have:

$$\begin{aligned} \mu_0^{-\frac{q_1}{d_{01}}} \mu_1^{\frac{q_0}{d_{01}}} &= \mu_0^{(-\frac{q_1}{q_0} c^0_{12} - \frac{q_2 d_{01}}{q_0 d_{012}})} \mu_1^{c^0_{12}} \mu_2^{\frac{d_{01}}{d_{012}}} = \cdots = \\ &= \mu_0^{(-\frac{q_1}{q_0} c^0_{1n} - \frac{q_2}{q_0} c^0_{2n} - \cdots - \frac{q_n d_n}{q_0 d_{0 \cdots n}})} \mu_1^{c^0_{1n}} \cdots \mu_{n-1}^{c^0_{n-1,n}} \mu_n^{\frac{d_n}{d_{0 \cdots n}}} = 1 \end{aligned}$$

So

$$\begin{cases} \mu_0^{-\frac{q_1}{d_{01}}} = \mu_1^{\frac{q_0}{d_{01}}}, \\ \mu_0^{(\frac{q_1}{q_0}c_{12}^0+\frac{q_2 d_{01}}{q_0 d_{012}})} = \mu_1^{c_{12}^0}\mu_2^{\frac{d_{01}}{d_{012}}}; \\ \cdots\cdots\cdots\cdots\cdots\cdots \\ \mu_0^{(\frac{q_1}{q_0}c_{1n}^0+\frac{q_2}{q_0}c_{2n}^0+\cdots+\frac{q_n d_n}{q_0 d_{0\cdots n}})} = \mu_1^{c_{1n}^0}\cdots\mu_{n-1}^{c_{n-1,n}^0}\mu_n^{\frac{d_n}{d_{0\cdots n}}} \end{cases}$$

We have

$$\mu_0 = (\mu_0^{\frac{1}{q_0}})^{q_0}, \mu_1 = (\mu_0^{\frac{1}{q_0}})^{q_1}, \mu_2 = (\mu_0^{\frac{1}{q_0}})^{q_2}, \cdots, \mu_n = (\mu_0^{\frac{1}{q_0}})^{q_n}. \tag{3.2}$$

We obtain the following results. if $(\mu_0, \cdots, \mu_n) \in G \subset (\mathbf{C}^*)^{n+1}$, then $(\mu_0, \cdots, \mu_n)$ satisfies the condition (3.2). So $G \cong \mathbf{C}^*$, let $\mu_0^{\frac{1}{q_0}} = \lambda$, then the G action on $(\mathbf{C}^*)^{\mathbf{n+1}}$ by $\lambda(x_0, \cdots, x_n) = (\lambda^{q_0}x_0, \cdots, \lambda^{q_n}x_n)$. Thus $\mathbf{P}^n_{q_0,\cdots,q_n} = (\mathbf{C}^{n+1})^*/\mathbf{C}^*$.

Proposition 3.3.1. *Let $Q = (q_0, \cdots, q_n)$ and $gcd(q_0, \cdots, q_n) = 1$, then the definitions of $\mathbf{P}^n_{q_0,\cdots,q_n}$ in the sections 3.1 and 3.2 are equivalent.*

4. The Chen-Ruan Cohomology Groups of Weighted Projective Spaces

4.1. **The Ordinary Cohomology Groups of Weighted Projective Spaces.**

Let $Q = (q_0, \cdots, q_n)$, $gcd(q_0, \cdots, q_n) = 1$ and $\mathbf{P}^n_{q_0,\cdots,q_n}$ be the weighted projective space of the type Q. The ordinary cohomology group of $\mathbf{P}^n_{q_0,\cdots,q_n}$ has already been studied by several authors, [1], [17]. Here we only give the results.

Theorem 4.1.1. *Let $Q = (q_0, \cdots, q_n)$ and $\mathbf{P}^n_{q_0,\cdots,q_n}$ be the weighted projective space of the type Q, then the cohomology group of $\mathbf{P}^n_{q_0,\cdots,q_n}$ with rational coefficient is*

$$H^i(\mathbf{P}^n_{q_0,\cdots,q_n}, \mathbf{Q}) = \begin{cases} \mathbf{Q}, & \text{if } i = 2r, 0 \le r \le n; \\ 0, & \text{if } i \text{ is odd or } i > 2n. \end{cases}$$

4.2. **Orbiford Structure from Toric Varieties.**

Let $Q = (q_0, \cdots, q_n)$ and $\mathbf{P}^n_{q_0\cdots,q_n}$ be the weighted projective space of the type Q. And let $\Xi = \{v_0, \cdots, v_n\}$ be the fan of $\mathbf{P}^n_{q_0,\cdots,q_n}$ in the lattice $\mathbf{Z}^n$ defined in section 3.2. We have the following proposition.

Proposition 4.2.1. *Let $Q = (q_0, \cdots, q_n)$ and $\mathbf{P}^n_{q_0,\cdots,q_n}$ be the weighted projective space of the type Q. Then taking $Q = (q_0, \cdots, q_n)$ as toric variety and its fan Ξ is generated by $\{v_0, \cdots, v_n\}$ above, it has the orbifold structure as follows.*

$$\{(U_{\sigma'_k}, G_{\sigma_k}, \pi_{\sigma_k}) | \sigma_k = (v_0, \cdots, \hat{v}_k, \cdots, v_n), k = 0, ..., n\}$$

In particular, $G_{\sigma_k} = \mathbf{Z}_{q_k}$ is the cyclic group of order q_k.

Proof. Because the fan Ξ of toric variety $\mathbf{P}^n_{q_0,\cdots,q_n}$ is generated by $\{v_0, \cdots, v_n\}$, so we all have $n+1$ n-dimensional cones $\sigma_0, \cdots, \sigma_n$. From section 2.4, we can conclude that

$$\{(U_{\sigma'_k}, G_{\sigma_k}, \pi_{\sigma_k}) | \sigma_k = (v_0, \cdots, \hat{v}_k, \cdots, v_n), k = 0, ..., n\}$$

forms the orbifold structure of $\mathbf{P}^n_{q_0,\cdots,q_n}$. Now we prove that for any σ_k, $G_{\sigma_k} = \mathbf{Z}_{q_k}$. From section 2.4 we have $G_{\sigma_k} = N/N_{\sigma_k}$, let $\pi_k : \mathbf{Z} \longrightarrow \mathbf{Z}_{q_k}$ be the standard projection. Define the map $Q_k : \mathbf{Z}^n \longrightarrow \mathbf{Z}$ such that $Q_k(a_1, \cdots, a_n) = q_0 a_1 + \cdots + q_n a_n, Q_i = (q_0, \cdots, \hat{q_k}, \cdots, q_n)$. Then we let

$$N_{Q,q_k} := Ker(\mathbf{Z}^n \longrightarrow \mathbf{Z} \longrightarrow \mathbf{Z}_{q_k})$$

From H.Conrads [5], because $\Xi = \{v_0, \cdots, v_n\}$ is the fan of $\mathbf{P}^n_{q_0,\cdots,q_n}$, so N_{Q,q_k} is generated by $\sigma_k = (v_0, \cdots, \hat{v_k}, \cdots, v_n)$, i.e. $N_{Q,k} = N_{\sigma_k}$. So we have:

$$G_{\sigma_k} = N/N_{\sigma_k} = \mathbf{Z}^n/N_{Q,q_k} \cong \mathbf{Z}_{q_k}.$$

□

Remark 4.2.2. From the proposition above, $U_{\sigma_k'} = \mathbf{C}^n$, so $U_{\sigma_k} = U_{\sigma_k'}/G_{\sigma_k} = \mathbf{C}^n/\mathbf{Z}_{q_k}$, and the action of $\mathbf{Z}_{q_k}$ is the diagonal action, its matrix representation can be computed from the method of section 2.4 which we will use in the following section.

4.3. **Twisted Sectors of Weighted Projective Spaces and Degree Shifting Numbers.** Given a weighted projective space $\mathbf{P}(Q)$ of type $Q = (q_0, \cdots, q_n)$, we now discuss the twisted sectors of $\mathbf{P}(Q)$. First, taken as toric variety, $\mathbf{P}(Q)$ has the orbit decomposition $\mathbf{P}(Q) = \bigsqcup_{\tau \in \Xi} O_\tau$, we will determined O_τ. From section 2.4, if $\sigma \in \Xi$ is a n-dimensional cone, suppose $\sigma = (v_0, \cdots, v_{n-1})$, then $G_\sigma = \mathbf{Z}_{q_n}$. We also know that $G_\sigma = \{k_a = \Sigma_{i=0}^{n-1} a_i v_i : k_a \in N, a_i \in [0,1)\}$. If $\tau = (v_0, \cdots, v_{i-1})$ is a face of σ, then $G_\tau = \{g_a \in G_\sigma : a_i = 0 \text{ if } j+1 \leq i \leq n\}$, i.e. $G_\tau = \{k_a = \Sigma_{j=1}^{i-1} a_j v_j : k_a \in N, a_j \in [0,1)\}$. G_τ can be taken as the local group of the points in O_τ. For the weighted projective space, we have the following proposition.

Proposition 4.3.1. *Given a weighted projective space $\mathbf{P}(Q)$ of type $Q = (q_0, \cdots, q_n)$. Let $\Xi = \{v_0, \cdots, v_n\}$ be the fan of $\mathbf{P}(Q)$. If $\tau = (v_0, \cdots, v_{i-1})$is a cone of Ξ, then we have $G_\tau \cong \mathbf{Z}_d$, where $d = gcd(q_i, \cdots, q_n)$. In particular, $(q_i, \cdots, q_n)$ is the maximal subset of $(q_0, \cdots, q_n)$ whose gcd is d iff the dimension of fixed point set of $\mathbf{Z}_d$ is $n - i$.*

Proof. Let $\sigma_k = (v_0, \cdots, v_{i-1}, v_i, \cdots, \hat{v_k}, \cdots, v_n), i \leq k \leq n$, then from proposition 4.2.1, $G_{\sigma_k} \cong \mathbf{Z}_{q_k}$. Since we have $G_{\sigma_k} = \{k_a = \Sigma_{j \neq k} a_j v_j : k_a \in N, a_j \in [0,1), i \leq k \leq n\}$, $G_\tau = \{k_a = \Sigma_{j=1}^{i-1} a_j v_j : k_a \in N, a_j \in [0,1)\}$, so G_τ is a subgroup of $G_{\sigma_k} (i \leq k \leq n)$. Next we show that any element of a common subgroup of $G_{\sigma_k} (i \leq k \leq n)$ has the form of the elements in G_τ. For any element $g_a \in \mathbf{Z}_{q_k} (i \leq k \leq n)$. Without loss of generality, assume $g_a \in \mathbf{Z}_{q_i}, \mathbf{Z}_{q_{i+1}}$, and

$$(1) g_a = a_0 v_0 + \cdots + a_{i-1} v_{i-1} + a_{i+1} v_{i+1} + \cdots + a_n v_n$$

$$(2) g_a = a_0' v_0 + \cdots + a_i' v_i + a_{i+2}' v_{i+2} + \cdots + a_n' v_n$$

Since the action of the same g_a at one neighborhood of a point is the same, so for $j \neq i, i+1, a_j = a_j'$, $(1) - (2)$, we obtain $a_{i+1} v_{i+1} - a_i' v_i = 0$, and we have a contradiction, because the generators v_{i+1} and v_i of the fan are independent over $N_{\mathbf{R}}$. We have $g_a = a_0 v_0 + \cdots + a_{i-1} v_{i-1}$ and $g_a \in G_\tau$. So we prove $G_\tau = \mathbf{Z}_d, d = gcd(q_i, \cdots, q_n)$.

Now if $(q_i, \cdots, q_n)$ is the maximal subset of $(q_0, \cdots, q_n)$ that satisfies the condition $gcd(q_i, \cdots, q_n) = d$, while the dimension of fixed point set of $\mathbf{Z}_d$ is not $n - i$. Because the dimension of O_τ is $n - i$, then we have a generator $g_a \in \mathbf{Z}_d$ and g_a can be represented by $g_a = a_0 v_0 + \cdots + a_{i-1} v_{i-1}$. And we must have some $a_s = 0$ for

$s \leq i-1$. If we let $\rho = (v_0, \cdots, \hat{v_s}, \cdots, v_{i-1})$, then from the first part of the proposition we have $G_\rho = \mathbf{Z}_d = gcd(q_s, q_i, \cdots, q_n)$ which will contradict the maximal principle of the condition.

Conversely, suppose the dimension of the fixed point set of $\mathbf{Z}_d$ is $n-i$, i.e. the dimension of the orbit O_τ. If we have a subset of $(q_0, \cdots, q_n)$ whose great common divisor is d, and the number of the subset is more than the $(q_i, \cdots, q_n)$. Without loss of generality, we assume $gcd(q_s, q_i, \cdots, q_n) = d$ for $0 \leq s \leq i-1$. Then let $\delta = (v_0, \cdots, v_{s-1}, v_{s+1}, \cdots, v_{i-1})$. From the first part of the theorem we have

$$G_\delta = \mathbf{Z}_d = \{k_a = a_0 v_0 + \cdots + a_{s-1}v_{s-1} + a_{s+1}v_{s+1} + \cdots + a_{i-1}v_{i-1} : a_i \in [0,1)\}$$

We see that the dimension of the fixed point set of $\mathbf{Z}_d$ overdues $n-i+1$, a contradiction. $\square$

Now we discuss the twisted sectors of the weighted projective space $\mathbf{P}(Q)$. From the theorem of Poddar in [21] for the twisted sectors of general toric varieties, we have:

Theorem 4.3.2. *([21]) A twisted sector of a weighted projective space is isomorphic to a subvariety $\overline{O}_\tau$ of $X_\Xi = \mathbf{P}(Q)$ for some $\tau \in \Xi$. There is a one to one correspondence between the set of twisted sectors of the type $\overline{O}_\tau$ and the set of integral vectors in the interior of τ which are linear combinations of the 1-dimensional generators of τ with coefficients in* $(0,1)$.

For the orbit O_τ of the weighted projective space $\mathbf{P}(Q)$ for some $\tau \in \Xi$, we have the following proposition:

Proposition 4.3.3. *Given a weighted projective space $\mathbf{P}(Q)$ of type $Q = (q_0, \cdots, q_n)$. Let $\Xi = \{v_0, \cdots, v_n\}$ be the fan of $\mathbf{P}(Q)$. If $\tau = (v_0, \cdots, v_{i-1})$ is a cone in Ξ, then $\overline{O}_\tau$ is the weighted projective space $\mathbf{P}(Q_\tau)$, where $Q_\tau = (q_i, \cdots, q_n)$.*

Proof. From Fulton [13], $\overline{O}_\tau$ is a toric variety and the fan $Star(\tau)$ can be described as follows. Let N_τ be the sublattice of N generated by $\tau \in N$, and $N(\tau) = N/N_\tau, M(\tau) = \tau^\perp \cap M$ be the quotient lattice and the dual. The star of a cone τ can be defined abstractly as the set of cones σ in Ξ that contain τ as a face. Such cones σ are determined by their images in $N(\tau)$, i.e. by

$$\overline{\sigma} = (\sigma + (N_\tau)_{\mathbf{R}})/(N_\tau)_{\mathbf{R}} \subset N_{\mathbf{R}}/(N_\tau)_{\mathbf{R}} = N(\tau)_{\mathbf{R}}$$

These cones $\{\overline{\sigma} : \tau < \sigma\}$ form a fan in $N(\tau)$, and we denote this fan by $Star(\tau)$, the corresponding toric variety is $n-k$-dimensional toric variety. For the toric variety $\mathbf{P}(Q)$, let $\overline{v}_i, \cdots, \overline{v}_n$ be the images of $v_0, \cdots, v_n$ in the quotient lattice $N(\tau)$, since in N, $q_0 v_0 + \cdots + q_n v_n = 0$, we have $q_i \overline{v}_i + \cdots + q_n \overline{v}_n = 0$ in the quotient lattice $N(\tau)$. So from the definition of weighted projective space, Remark 3.2.4, we conclude that the toric variety corresponding to the fan $Star(\tau)$ in the quotient lattice $N(\tau)$ is the weighted projective space $\mathbf{P}(Q_\tau)$, where $Q_\tau = (q_i, \cdots, q_n)$. $\square$

Remark 4.3.4. The gcd of $(q_i, \cdots, q_n)$ need not necessary be 1, if $d = gcd(q_i, \cdots, q_n)$, in general, $\mathbf{P}(Q_\tau)$ is a nonreduced orbifold, and it has a corresponding reduced orbifold $\mathbf{P}(Q_\tau^{'})$, where $Q_\tau^{'} = (q_i/d, \cdots, q_n/d)$.

Theorem 4.3.5. *Let $Q = (q_0, \cdots, q_n)$ and $\mathbf{P}(Q) = \mathbf{P}^n_{q_0, \cdots, q_n}$ be the weighted projective space of the type Q. Let $\Xi = \{v_0, \cdots, v_n\}$ be the fan of $\mathbf{P}(Q)$. If $\tau = (v_0, \cdots, v_{i-1})$ is a cone in Ξ, then $\overline{O}_\tau = \mathbf{P}(Q_\tau)$, where $Q_\tau = (q_i, \cdots, q_n)$. Let*

$gcd(q_i,\cdots,q_n)=d$. Then we have: $\overline{O}_\tau=\mathbf{P}(Q_\tau)$ is a twisted sector iff $(q_i,\cdots,q_n)$ is the maximal subset of $(q_0,\cdots,q_n)$ that satisfies the condition $gcd(q_i,\cdots,q_n)=d$.

Proof. If $\overline{O}_\tau$ is a twisted sector, then suppose $\overline{O}_\tau = X_{(g_a)}$, where $g_a\in G_\tau=\mathbf{Z}_d$ is a generator. If we have $q_k\bar{\in}(q_i,\cdots,q_n)$, $gcd(q_k,q_i,\cdots,q_n)=d$, let $\delta=(v_0,\cdots,\hat{v_k},\cdots,v_{i-1})$, then $\overline{O}_\delta=\mathbf{P}(Q_\delta)$, where $Q_\delta=(q_k,q_i,\cdots,q_n)$. Since from section 2.4, $G_\delta=\mathbf{Z}_d=\{k_a=\Sigma_{j=0}^{i-1}a_jv_j : j\neq k, a_j\in[0,1)\}$, the coefficients of the representation of g_a are non zeroes, so from theorem 4.3.2, $X_{(g_a)}=\overline{O}_\delta$ which contradicts the condition.

Suppose $(q_i,\cdots,q_n)$ is the maximal subset of $(q_0,\cdots,q_n)$ that meets the condition $gcd(q_i,\cdots,q_n)=d$, from proposition 4.4, let $\tau=(v_0,\cdots,v_{i-1}), G_\tau\cong\mathbf{Z}_d$ and the dimension of the fixed point set of $\mathbf{Z}_d$ is $n-i$, the dimension of the orbit O_τ. Since

$$G_\tau=\mathbf{Z}_d=\{k_a=\Sigma_{j=0}^{i-1}a_jv_j : a_j\in[0,1)\}$$

we must have one generator $g_a\in G_\tau$ so that g_a can be represented by

$$g_a=a_0v_0+\cdots+a_{i-1}v_{i-1}, \text{all } a_j\neq 0, j=0,\cdots,i-1.$$

Thus the dimension of the fixed point set of g_a is $n-i$. From theorem 4.3.2, we see that $X_{(g_a)}=\bar{O}_\tau$. □

Remark 4.3.6. From above analysis, if $\tau=(v_0,\cdots,v_{i-1})$ is a cone in Ξ, we describe the orbifold structure of twisted sector $X_{(g_a)}=\bar{O}_\tau$ as follows. In the points of O_τ, the local group at the point x is $\mathbf{Z}_d$, $d=gcd(q_i,\cdots,q_n)$, and $C(g_a)=\mathbf{Z}_d$ acts trivially. If $\delta>\tau$ is a cone, then $O_\delta\subset\overline{O}_\tau$,the local group at the $y\in O_\delta$ is the cyclic group $\mathbf{Z}_t$, where $t=gcd(q_{t_1},\cdots,q_{t_s}),q_i,\cdots,q_n$, and $\delta=(v_0,\cdots,\hat{v}_{t_1},\cdots,\hat{v}_{t_s},\cdots,v_{i-1})$. And the action of $\mathbf{Z}_t$ on $\bar{O}_\tau$ can be described as before.

The degree shifting numbers can be computed easily. For instance, let $X_{(g_a)}=\overline{O}_\tau$ be a twisted sector, and $\tau=(v_0,\cdots,v_{i-1})$, we can write g_a as

$$g_a=\Sigma_{j=0}^{i-1}a_jv_j, a_j\in(0,1)$$

The degree shifting number $\iota_{(g_a)}$ of $X_{(g_a)}$ is:

$$\iota_{(g_a)}=\Sigma_{j=0}^{i-1}a_j$$

So in the orbifold structure of weighted projective space $\mathbf{P}(Q)$, we can compute the degree shifting number corresponding to any twisted sector.

4.4. The Chen-Ruan Cohomology Groups of Weighted Projective Spaces.

Up to now, given a weighted projective space $\mathbf{P}(Q)$ of type $Q=(q_0,\cdots,q_n)$, from Theorem 4.3.5 we can calculate the twisted sectors of $\mathbf{P}(Q)$ and we also can compute the degree shifting numbers of the corresponding twisted sectors. So we write the Chen-Ruan cohomology group of $\mathbf{P}(Q)$ in the following manner.

Theorem 4.4.1. *Let $Q=(q_0,\cdots,q_n)$ and $\mathbf{P}(Q)=\mathbf{P}^n_{q_0,\cdots,q_n}$ be the weighted projective space of the type Q. Let $\Xi=\{v_0,\cdots,v_n\}$ be the fan of $\mathbf{P}(Q)$. Then the orbifold cohomology group of $\mathbf{P}(Q)$ is*

$$H^p_{orb}(\mathbf{P}(Q);\mathbf{Q})\cong\bigoplus_{\sigma\in\Xi,l\in\mathbf{Q}}H^{p-2l}(\overline{O}_\sigma)\otimes\oplus_{t\in\sigma_l}\mathbf{Q}t$$

Where $\sigma_l = \{\sum_{v_i \subset \sigma} a_i v_i \in N : a_i \in (0,1), \sum_{v_i \subset \sigma} a_i = l\}$.*(when* $\sigma = 0$*, set* $l = 0, T(\sigma)_l = \mathbf{C}$*).* $\overline{O}_\sigma$ *is the closure of the orbit corresponding to* $\sigma \in \Xi$*. Here* p *is rational numbers in* $[0, n]$*, and* $H^{p-2l}(\overline{O}_\sigma) = 0$ *if* $p - 2l$ *is not integral.*

Note that the elements of $\oplus_{0 \neq \sigma \in \Delta, l} \sigma_l$ correspond to the twisted sectors of $\mathbf{P}(Q)$.

4.5. **Example.** For $Q = (2,3,4)$, $\mathbf{P}(Q) = \mathbf{P}^2_{2,3,4}$, we have $q_0 = 2, q_1 = 3, q_2 = 4$. From proposition 3.2.2., we have

$$C_0 = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$$

So let $v_1 = (2,0), v_2 = (0,1)$, we have $v_0 = -\frac{q_1}{q_0} v_1 - \frac{q_2}{q_0} v_2 = (-3,-2)$. The fan Ξ of $\mathbf{P}^2_{2,3,4}$ is generated by $\{v_0, v_1, v_2\}$. For $\sigma_2 = (v_0, v_1) = ((-3,-2),(2,0))$, we have $G_{\sigma_2} = N/N_{\sigma_2} = \mathbf{Z}_4$. We write the matrix representation of the action of $\mathbf{Z}_4$ on $U_{\sigma_2'} = \mathbf{C}^2$ as follows:

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \begin{pmatrix} e^{2\pi i \cdot \frac{1}{2}} & 0 \\ 0 & e^{2\pi i \cdot \frac{1}{4}} \end{pmatrix}; \begin{pmatrix} 1 & 0 \\ 0 & e^{2\pi i \cdot \frac{1}{2}} \end{pmatrix}; \begin{pmatrix} e^{2\pi i \cdot \frac{1}{2}} & 0 \\ 0 & e^{2\pi i \cdot \frac{3}{4}} \end{pmatrix}$$

For $\sigma_1 = (v_0, v_2) = ((-3,-2),(0,1))$, we have $G_{\sigma_1} = \mathbf{Z}_3$. We write the matrix representation of the action of $\mathbf{Z}_3$ on $U_{\sigma_1'} = \mathbf{C}^2$ as follows:

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \begin{pmatrix} e^{2\pi i \cdot \frac{1}{3}} & 0 \\ 0 & e^{2\pi i \cdot \frac{2}{3}} \end{pmatrix}; \begin{pmatrix} e^{2\pi i \cdot \frac{2}{3}} & 0 \\ 0 & e^{2\pi i \cdot \frac{1}{3}} \end{pmatrix}$$

And for $\sigma_0 = (v_1, v_2) = ((2,0),(0,1))$, we have $G_{\sigma_0} = \mathbf{Z}_2$ and representation:

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \begin{pmatrix} e^{2\pi i \cdot \frac{1}{2}} & 0 \\ 0 & 1 \end{pmatrix}$$

If we let

$$g_1 = \begin{pmatrix} e^{2\pi i \cdot \frac{1}{2}} & 0 \\ 0 & e^{2\pi i \cdot \frac{1}{4}} \end{pmatrix}; g_2 = \begin{pmatrix} e^{2\pi i \cdot \frac{1}{3}} & 0 \\ 0 & e^{2\pi i \cdot \frac{2}{3}} \end{pmatrix}; g_3 = \begin{pmatrix} e^{2\pi i \cdot \frac{1}{2}} & 0 \\ 0 & 1 \end{pmatrix}$$

then we have the twisted sectors: let $p_1 = [0,1,0], p_2 = [0,0,1]$, $X_{(g_1)} = X_{(g_1^3)} = p_2, \iota_{(g_1)} = \frac{3}{4}, \iota_{(g_1^3)} = \frac{5}{4}$; $X_{(g_1^2)}$ and $X_{(g_3)}$ are the same twisted sector $\mathbf{P}_{2,0,4}$, and $\iota_{(g_1^2)} = \iota_{(g_3)} = \frac{1}{2}$; $X_{(g_2)} = X_{(g_2^2)} = p_1, \iota_{(g_2)} = \iota_{(g_2^2)} = 1$. So

$$\begin{aligned} H^p_{orb}(\mathbf{P}^2_{2,3,4}; \mathbf{Q}) &= H^p(\mathbf{P}^2_{2,3,4}; \mathbf{Q}) \oplus H^{p-2\iota_{(g_1)}}(\{p_2\}; \mathbf{Q}) \oplus H^{p-2\iota_{(g_1^3)}}(\{p_2\}; \mathbf{Q}) \\ &\oplus H^{p-2\iota_{(g_1^2)}}(\mathbf{P}_{2,0,4}; \mathbf{Q}) \oplus 2H^{p-2\iota_{(g_2)}}(\{p_1\}; \mathbf{Q}) \end{aligned}$$

We compute the orbifold cohomology group of $\mathbf{P}^2_{2,3,4}$ as

$$\begin{aligned} &H^0_{orb}(\mathbf{P}^2_{2,3,4}; \mathbf{Q}) = \mathbf{Q}; \\ &H^1_{orb}(\mathbf{P}^2_{2,3,4}; \mathbf{Q}) = \mathbf{Q}; \\ &H^{\frac{3}{2}}_{orb}(\mathbf{P}^2_{2,3,4}; \mathbf{Q}) = \mathbf{Q}; \\ &H^2_{orb}(\mathbf{P}^2_{2,3,4}; \mathbf{Q}) = \mathbf{Q} \oplus \mathbf{Q} \oplus \mathbf{Q}; \\ &H^{\frac{5}{2}}_{orb}(\mathbf{P}^2_{2,3,4}; \mathbf{Q}) = \mathbf{Q}; \\ &H^3_{orb}(\mathbf{P}^2_{2,3,4}; \mathbf{Q}) = \mathbf{Q}; \\ &H^4_{orb}(\mathbf{P}^2_{2,3,4}; \mathbf{Q}) = \mathbf{Q}. \end{aligned}$$

All the other dimensions of the Chen-Ruan cohomology groups are zero.

5. The Chen-Ruan Cohomology Rings of the Weighted Projective Spaces

5.1. **The Ordinary Cohomology Ring of Weighted Projective Spaces.** In this section we recall the ordinary cohomology ring of the weighted projective space. The readers may refer to [1]. Let $Q = (q_0, \cdots, q_n)$ and $\mathbf{P}(Q) = \mathbf{P}^n_{q_0,\cdots,q_n}$ be the weighted projective space of the type Q. Let $\mathbf{P}^n$ be the n-dimensional complex projective space. As in [1], let $\varphi : \mathbf{P}^n \longrightarrow \mathbf{P}(Q)$ be the map taking $[x_0, \cdots, x_n]$ to $[x_0^{q_0}, \cdots, x_n^{q_n}]$. Take $k \in \{0, \cdots, n\}$, and consider $I = \{i_0, \cdots, i_k\}$ with $0 \le i_0 < \cdots < i_k \le n$. Put $l_I = l_I(q_{i_0}, \cdots, q_{i_k}) = q_{i_0} \cdots q_{i_k}/gcd(q_{i_0}, \cdots, q_{i_k})$, and let

$$l_k = l_k(q_0, \cdots, q_n) = lcm\{l_I | I \subset \{0, \cdots, n\}, |I| = k + 1\}$$

Theorem 5.1.1. *([1]) For each $k, 0 \le k \le n$, there exists a unique $\xi_k \in H^{2k}(\mathbf{P}(Q); \mathbf{Q})$ such that $\varphi^*(\xi_k) = l_k \xi^k$, and $\{1, \xi, \cdots, \xi^n\}$ is a $\mathbf{Q}$-basis of the free abelian group $H^{2k}(\mathbf{P}(Q); \mathbf{Q})$. In other words there are commutative diagrams*

$$\begin{tikzcd}
H^{2k}(\mathbf{P}(Q);\mathbf{Q}) \arrow[r, "\varphi^*"] \arrow[d, equal] & H^{2k}(\mathbf{P}(Q);\mathbf{Q}) \arrow[d, equal] \\
\mathbf{Q} \arrow[r, "\cdot l_k"'] & \mathbf{Q}
\end{tikzcd}$$

So we can make precise the multiplicative structure of the cohomology $H^{2k}(\mathbf{P}(Q); \mathbf{Q})$. Since $\varphi^* : H^*(\mathbf{P}(Q); \mathbf{Q}) \longrightarrow H^*(\mathbf{P}^n; \mathbf{Q})$ is a ring homomorphism, so

$$\xi_i \xi_j = \begin{cases} e_{ij}\xi_{i+j}, & i + j \le n; \\ 0, & \text{if not.} \end{cases}$$

Where $e_{ij} = l_i l_j / l_{i+j}, 1 \le i, j \le n$.

In the polynomial ring $\mathbf{Q}[T_1, \cdots, T_n]$, let A be the idea generated by the elements

$$T_i T_j (i + j > n) \text{ and } T_i T_j - e_{ij} T_{i+j} (i + j \le n)$$

We obtain a ring isomorphism

$$H^*(\mathbf{P}(Q); \mathbf{Q}) \cong \mathbf{Q}[T_1, \cdots, T_n]/A$$

Where ξ_i corresponds to the class of T_i.

5.2. **Three Multi-sectors.** Let $\mathbf{P}(Q)$ be the weighted projective space of type $Q = (q_0, \cdots, q_n)$, and $\Xi = \{v_0, \cdots, v_n\}$ be the fan when $\mathbf{P}(Q)$ is taken as toric variety. Denote $\Xi[n]$ the set of n-dimensional cones. For a cone $\tau \in \Xi$, denote the set of its primitive 1-dimensional generators by $\tau[1]$. From the section 2.4, $G_\tau = \{\Sigma_{v_i \subset \tau[1]} a_i v_i : a_i \in [0, 1)\}$, let $R(\tau) := \{g_a = \Sigma a_i u_i | u_i \in \tau[1], 0 \le a_i < 1\} \cap N$. Now we describe the 3-multisector $X_{(\mathbf{g})}$ for $\mathbf{g} = (g_1, g_2, g_3)$.

Take any $x \in X_\Xi = \mathbf{P}(Q)$ with nontrivial local group. Then x belongs to a unique O_τ such that τ is not the trivial cone. Pick any elements g_a, g_b from $G_x = G_\tau$. We shall find $X_{(\mathbf{g})}$ where $\mathbf{g} = (g_a, g_b, (g_a g_b)^{-1})$. Let τ_a and τ_b be the faces of τ, whose interiors contain g_a and g_b respectively. Let σ be any n-dimensional cone containing τ. Let z be any point in $U_{\sigma'}$. Suppose z is fixed by both g_a and g_b. Then $\chi^{u^i} = 0$ whenever $u_i \in \tau_a \cup \tau_b$. Hence $\pi_\sigma(z) \in \overline{O}_{\tau_a} \cap \overline{O}_{\tau_b} \cap U_\sigma$. A local uniformizing system for $X_{(\mathbf{g})}$ is given by $(V_x^{\mathbf{g}}, G_x, \pi)$, where

$$V_x^{\mathbf{g}} = (\mathbf{C}^j \times W) \cap \{\chi^{u_i} = 0 : \forall u_i \in \tau_a \cup \tau_b\}$$

This leads us to observe that $\{(x,\mathbf{g}) \in X_{(\mathbf{g})} | x \in U_\sigma\}$ is complex analytical isomorphic to $\overline{O}_{\tau_a} \cap \overline{O}_{\tau_b} \cap U_\sigma$. Since this is true in respective of the choice of σ, $X_{(\mathbf{g})} \cong \overline{O}_{\tau_a} \cap \overline{O}_{\tau_b}$. So we have:

Theorem 5.2.1. *([22]) If $\tau_1[1] \cup \tau_2[1]$ generates an element of Ξ, then for every pair $g_{a_1} \in R(\tau_1) \cap Int(\tau_1), g_{a_2} \in R(\tau_2) \cap Int(\tau_2)$, we have a unique 3-multisector $X_{(\mathbf{g})}$, which is analytically isomorphic to $\overline{O}_{\tau_1} \cap \overline{O}_{\tau_2}$. As we vary over τ_1, τ_2, we obtain all the 3-multisectors.*

Since the fan $\Xi = \{v_0, \cdots, v_n\}$ of $\mathbf{P}(Q)$ for $Q = (q_0, \cdots, q_n)$ has $n+1$ primitive 1-dimensional generators. If τ_1 and τ_2 are two cones of Ξ, then we have that $\tau_1[1] \cup \tau_2[1]$ form an element $\tau = \tau_1[1] \cup \tau_2[1]$ of Ξ, so $\overline{O}_{\tau_1} \cap \overline{O}_{\tau_2}$ is an 3-multisector of $\mathbf{P}(Q)$. Moreover, we can prove the 3-multisectors of $\mathbf{P}(Q)$ are actually twisted sectors.

Theorem 5.2.2. *Let $X = \mathbf{P}(Q)$ be the weighted projective space of type $Q = (q_0, \cdots, q_n)$. $X_{(g_1)}$ and $X_{(g_2)}$ are two twisted sectors of X, suppose they correspond to the cones τ_1 and τ_2 respectively, i.e. $X_{(g_1)} = \overline{O}_{\tau_1}, X_{(g_2)} = \overline{O}_{\tau_2}$. Then $X_{(\mathbf{g})} = X_{(g_1, g_2, (g_1 g_2)^{-1})}$ is still a twisted sector.*

Proof. First if $\tau_1 \subset \tau_2$, then from theorem 5.2.1, $\tau_1[1] \cup \tau_2[1]$ generates τ_2, so $X_{(\mathbf{g})} = \overline{O}_{\tau_2} = X_{(g_2)}$.

If $\tau_1[1] \cap \tau_2[1] = \emptyset$, let $\tau = (\tau_1[1] \cup \tau_2[1])$, then $X_{(\mathbf{g})} = X_{(g_1, g_2, (g_1 g_2)^{-1})} = \overline{O}_\tau$. Since $G_\tau = \{\Sigma_{v_i \subset \tau} a_i v_i | a_i \in [0,1)\}$, we always can find an element $g \in G_\tau$ such that $g = \Sigma_{v_i \subset \tau} a_i v_i$, all $a_i \neq 0$. This follows if we take $g_1 = \Sigma_{v_i \subset \tau_1} a_i v_i (a_i \neq 0)$ and $g_2 = \Sigma_{v_i \subset \tau_2} a_i v_i (a_i \neq 0)$. From theorem 4.3.2, $X_{(\mathbf{g})} = \overline{O}_\tau = X_{(g)}$.

If τ_1 and τ_2 do not satisfy the above two types of conditions, without loss of generality, we suppose

$$\tau_1 = (v_0, \cdots, v_s), \tau_2 = (v_0, \cdots, v_j, v_{s+1}, \cdots, v_t), j < s, t > s.$$

then let $\tau = \tau_1[1] \cup \tau_2[1] = (v_0, \cdots, v_j, \cdots, v_s, v_{s+1}, \cdots, v_t)$. From proposition 4.3.3, we know $\overline{O}_\tau = \mathbf{P}(Q_\tau)$, where $Q_\tau = (0, \cdots, 0, q_{t+1}, \cdots, q_n)$. While $\overline{O}_{\tau_1} = \mathbf{P}(Q_{\tau_1})$, where $Q_{\tau_1} = (0, \cdots, 0, q_{s+1}, \cdots, q_t, q_{t+1}, \cdots, q_n)$, $\overline{O}_{\tau_2} = \mathbf{P}(Q_{\tau_2})$, where $Q_{\tau_2} = (0, \cdots, 0, q_{j+1}, \cdots, q_s, 0, \cdots, 0, q_{t+1}, \cdots, q_n)$. Let

$$d_1 = gcd(q_{s+1, \cdots, q_n}), \ d_2 = gcd(q_{j+1}, \cdots, q_s, q_{t+1}, \cdots, q_n) \tag{5.1}$$

So from theorem 4.3.5, $(q_{s+1}, \cdots, q_n)$ and $(q_{j+1}, \cdots, q_s, q_{t+1}, \cdots, q_n)$ are the maximal subsets of $(q_0, \cdots, q_n)$ that satisfy the condition (5.1). We conclude that $gcd(q_{t+1}, \cdots, q_n) \geq d_1 d_2$ and that $(q_{t+1}, \cdots, q_n)$ must be the maximal subset of $(q_0, \cdots, q_n)$ that satisfies this condition. So from theorem 4.3.5., $\overline{O}_\tau$ is a twisted sector. □

Remark 5.2.3. From the above theorem, every 3-multisector of weighted projective space $\mathbf{P}(Q)$ of type $Q = (q_0, \cdots, q_n)$ is actually a twisted sector, so we can refer to the Remark 4.3.6. to describe the orbifold structure of the 3-multisectors.

5.3. **The Chen-Ruan Cohomology Rings of Weighted Projective Spaces.** In this section we discuss the key point of computing the ring structure of Chen-Ruan cohomology of weighted projective space $\mathbf{P}(Q)$ of type $Q = (q_0, \cdots, q_n)$, i.e., the orbifold cup product. The most important part for the orbifold cup product is the obstruction bundle which was constructed as follows.

Let $X_{(\mathbf{g})}$ be a 3-multisector of $X = \mathbf{P}(Q)$, $\mathbf{g} = (g_1, g_2, g_3) \in T_3^0$. Let $E_{(\mathbf{g})} \longrightarrow X_{(\mathbf{g})}$ be the obstruction bundle defined in the section 2.3.1. On the local chart $(V_x^{\mathbf{g}}, C(\mathbf{g}), \pi)$ of $X_{(\mathbf{g})}$, $E_{(\mathbf{g})}$ is given by $(TV_x \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})} \times V_x^{\mathbf{g}} \longrightarrow V_x^{\mathbf{g}}$, and the rank of $E_{(\mathbf{g})}$ is given by the formula(1.3).

$$rank_{\mathbf{C}}(E_{(\mathbf{g})}) = dim_{\mathbf{C}}(X_{(\mathbf{g})}) - dim_{\mathbf{C}}(X) + \Sigma_{j=1}^{3} \iota_{(g_j)} \tag{5.2}$$

The orbifold cup product is defined by: let $\eta_j \in H^{d_j}(X_{(g_j)}; \mathbf{Q})$, for $j = 1, 2, 3$. Define maps $e_j : X_{(\mathbf{g})} \longrightarrow X_{(g_j)}$ by $(x, (\mathbf{g})_x) \longmapsto (p, (g_j)_x)$. Then

$$< \eta_1, \eta_2, \eta_3 >_{orb} = \int_{X_{(\mathbf{g})}}^{orb} e_1^* \eta_1 \wedge e_2^* \eta_2 \wedge e_3^* \eta_3 \wedge e_A(E_{(\mathbf{g})}) \tag{5.3}$$

Where $e_A(E_{(\mathbf{g})})$ is the Euler form computed from the connection A.

$$< \eta_1 \cup_{orb} \eta_2, \eta_3 >_{orb} = < \eta_1, \eta_2, \eta_3 >_{orb} \tag{5.4}$$

From the above, we have the following proposition.

Proposition 5.3.1. *Let* $\alpha \in H^*_{orb}(X_{(g_1)}; \mathbf{Q})$, $\beta \in H^*_{orb}(X_{(g_2)}; \mathbf{Q})$, $\mathbf{g} = (g_1, g_2, g_3) \in T_3^0$, *if* $\Sigma_{j=1}^3 \iota_{(g_j)} > n$, *then* $\alpha \cup_{orb} \beta = 0$.

Proof. From (5.2), we have $\Sigma_{j=1}^3 \iota_{(g_j)} - n = rank_{\mathbf{C}}(E_{(\mathbf{g})}) - dim_{\mathbf{C}}(X_{(\mathbf{g})})$. If $\Sigma_{j=1}^3 \iota_{(g_j)} > n$, then $rank_{\mathbf{C}}(E_{(\mathbf{g})}) > dim_{\mathbf{C}}(X_{(\mathbf{g})})$, so the integration (5.3) is zero, $\alpha \cup_{orb} \beta = 0$. □

Now in the next three sections we concretely discuss how to compute the 3-point function defined in (5.3).

5.4. $\mathbf{q_0}, \cdots, \mathbf{q_n}$ **are mutually prime-A Simple Case.** Given a $n+1$-tuple $Q = (q_0, \cdots, q_n)$. $q_i's$ are mutually prime. Let $\mathbf{P}(Q)$ be the weighted projective space of type Q, then the orbifold singularities are the $n+1$ isolated points: $p_i = [0, \cdots, i, \cdots 0] (i = 0, 1, \cdots, n)$ with local orbifold groups $\mathbf{Z}_{q_i} (i = 0, 1, \cdots, n)$. If we let $c_0, \cdots, c_n$ be the generators of $\mathbf{Z}_{q_0}, \cdots, \mathbf{Z}_{q_n}$ respectively, then we have $q_0 - 1$ twisted sectors isomorphic to $X_{(c_0)} = p_0$,$\cdots$,$q_n - 1$ twisted sectors isomorphic to $X_{(c_n)} = p_n$. And we can also see that the 3-sectors are all isolated points.

If we have $\alpha \in H^*(X_{(g_1)}; \mathbf{Q})$,$\beta \in H^*(X_{(g_2)}; \mathbf{Q})$, then $X_{(g_1, g_2, (g_1 g_2)^{-1})} = \{pt\}$ iff g_1, g_2 belong to some $\mathbf{Z}_{q_i} (i = 0, 1, \cdots, n)$. Without loss of generality, we assume $g_1, g_2 \in \mathbf{Z}_0$, then from the formula (4.1.7) in [6],

$$\alpha \cup_{orb} \beta = \sum_{(h_1, h_2), h_i \in (g_i)} (\alpha \cup_{orb} \beta)_{(h_1, h_2)} \tag{5.5}$$

where

$$< (\alpha \cup_{orb} \beta)_{(h_1, h_2)}, \gamma >_{orb} = \int_{X_{(h_1, h_2)}} e_1^* \alpha \wedge e_2^* \beta \wedge e_3^* \gamma \wedge e(E_{(\mathbf{g})}) \tag{5.6}$$

$e_i : X_{(\mathbf{g})} \longrightarrow X_{(g_i)}$ is the map mentioned above and $E_{(\mathbf{g})}$ is the obstruction bundle over $X_{(\mathbf{g})}$. From the formula (4.2), the dimension of the bundle $E_{(\mathbf{g})}$ is:

$$dim(e(E_{(\mathbf{g})})) = 2(\iota_{(g_1)} + \iota_{(g_2)} + \iota_{(g_3)}) - 2n.$$

Because $X_{(\mathbf{g})}$ is a point, so the integration (5.6) is nonzero iff $\alpha \in H^0(X_{(g_1)}; \mathbf{Q})$,$\beta \in H^0(X_{(g_2)}; \mathbf{Q})$,$\gamma \in H^0(X_{(g_3)}; \mathbf{Q})$ and $dim(e(E_{(\mathbf{g})})) = 0$. At this moment $\iota_{(g_1)} +$

$\iota_{(g_2)} + \iota_{(g_3)} = n$. Suppose α and β are all generators. Let $\gamma \in H^0(X_{(g_3)}; \mathbf{Q})$ is the generator. So the integration (5.6) is:

$$< (\alpha \cup_{orb} \beta)_{(g_1,g_2)}, \gamma >_{orb} = \frac{1}{|\mathbf{Z}_{q_0}|} \int_{\{pt\}} e_1^*\alpha \wedge e_2^*\beta \wedge e_3^*\gamma = \frac{1}{q_0}$$

If we let the δ be the generator of $H^0(X_{(g_1 g_2)}; \mathbf{Q})$, then the integration

$$< \delta, \gamma >_{orb} = \int_{X_{(g_1 g_2)}} \delta \wedge I^*\gamma = \frac{1}{q_0}$$

So we have

$$\alpha \cup_{orb} \beta = \delta. \tag{5.7}$$

Example 5.4.1. For $Q = (2,3,5)$, $\mathbf{P}(Q) = \mathbf{P}^2_{2,3,5}$, we have $q_0 = 2, q_1 = 3, q_2 = 5$. From proposition 3.2.2., we have

$$C_0 = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}$$

So let $v_1 = (2,1), v_2 = (0,1)$, we have $v_0 = -\frac{q_1}{q_0}v_1 - \frac{q_2}{q_0}v_2 = (-3,-4)$. The fan of $\mathbf{P}^2_{2,3,4}$ is generated by $\{v_0, v_1, v_2\}$. For $\sigma_2 = (v_0, v_1) = ((-3,-4),(2,1))$, we have $G_{\sigma_2} = N/N_{\sigma_2} = \mathbf{Z}_5$. We write the matrix representation of the action of $\mathbf{Z}_5$ on $U_{\sigma_2'} = \mathbf{C}^2$ as follows.

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \begin{pmatrix} e^{2\pi i\cdot\frac{1}{5}} & 0 \\ 0 & e^{2\pi i\cdot\frac{4}{5}} \end{pmatrix}; \begin{pmatrix} e^{2\pi i\cdot\frac{2}{5}} & 0 \\ 0 & e^{2\pi i\cdot\frac{3}{5}} \end{pmatrix}$$

$$\begin{pmatrix} e^{2\pi i\cdot\frac{3}{5}} & 0 \\ 0 & e^{2\pi i\cdot\frac{2}{5}} \end{pmatrix}; \begin{pmatrix} e^{2\pi i\cdot\frac{4}{5}} & 0 \\ 0 & e^{2\pi i\cdot\frac{1}{5}} \end{pmatrix}$$

For $\sigma_1 = ((-3,-4),(0,1))$, we have $G_{\sigma_1} = \mathbf{Z}_3$. We write the matrix representation of the action of $\mathbf{Z}_3$ on $U_{\sigma_1'} = \mathbf{C}^2$ as follows.

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \begin{pmatrix} e^{2\pi i\cdot\frac{1}{3}} & 0 \\ 0 & e^{2\pi i\cdot\frac{1}{3}} \end{pmatrix}; \begin{pmatrix} e^{2\pi i\cdot\frac{2}{3}} & 0 \\ 0 & e^{2\pi i\cdot\frac{2}{3}} \end{pmatrix}$$

And for $\sigma_0 = (v_1, v_2) = ((2,1),(0,1))$, we have $G_{\sigma_0} = \mathbf{Z}_2$ and representation:

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}; \begin{pmatrix} e^{2\pi i\cdot\frac{1}{2}} & 0 \\ 0 & e^{2\pi i\cdot\frac{1}{2}} \end{pmatrix}$$

If we let

$$g_1 = \begin{pmatrix} e^{2\pi i\cdot\frac{1}{5}} & 0 \\ 0 & e^{2\pi i\cdot\frac{4}{5}} \end{pmatrix}; g_2 = \begin{pmatrix} e^{2\pi i\cdot\frac{1}{3}} & 0 \\ 0 & e^{2\pi i\cdot\frac{1}{3}} \end{pmatrix}; g_3 = \begin{pmatrix} e^{2\pi i\cdot\frac{1}{2}} & 0 \\ 0 & e^{2\pi i\cdot\frac{1}{2}} \end{pmatrix}$$

then we have the twisted sectors: let $p_0 = [1,0,0], p_1 = [0,1,0], p_2 = [0,0,1]$, $X_{(g_1)} = X_{(g_1^2)} = X_{(g_1^3)} = X_{(g_1^4)} = p_0$, $X_{(g_2)} = X_{(g_2^2)} = p_1$ and $X_{(g_3)} = p_3$.

Since

$$\begin{aligned}
&\iota_{(g_1)} + \iota_{(g_1^4)} = 2, \\
&\iota_{(g_1^2)} + \iota_{(g_1^3)} = 2, \\
&\iota_{(g_2)} + \iota_{(g_2)} + \iota_{(g_2)} = 2, \\
&\iota_{(g_2)} + \iota_{(g_2^2)} = 2, \\
&\iota_{(g_3)} + \iota_{(g_3)} = 2.
\end{aligned}$$

Let $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ be the generators of $H^0(X_{(g_1)}; \mathbf{Q}), H^0(X_{(g_1^2)}; \mathbf{Q}), H^0(X_{(g_1^3)}; \mathbf{Q})$, $H^0(X_{(g_1^4)}; \mathbf{Q})$ respectively, β_1, β_2 be the generators of $H^0(X_{(g_2)}; \mathbf{Q}), H^0(X_{(g_2^2)}; \mathbf{Q})$, and γ be the generators of $H^0(X_{(g_3)}; \mathbf{Q})$. We also let e_0 be the generator of $H^0(\mathbf{P}^2_{2,3,5}; \mathbf{Q})$. So from the above discussion in section 5.4. and the formula (5.7), we have

$$\begin{aligned}
&\alpha_1 \cup_{orb} \alpha_4 = e_0, \\
&\alpha_2 \cup_{orb} \alpha_3 = e_0, \\
&\beta_1 \cup_{orb} \beta_1 = \beta_2, \\
&\beta_1 \cup_{orb} \beta_2 = e_0, \\
&\gamma \cup_{orb} \gamma = e_0.
\end{aligned}$$

5.5. **The Obstruction Bundle.** In this section we introduce a method to compute the obstruction bundle locally. Let $E_{(\mathbf{g}} \longrightarrow X_{(\mathbf{g})}$ be the obstruction bundle over the 3-multisector $X_{(\mathbf{g})}$. For a weighted projective space $\mathbf{P}(Q)$ of type $Q = (q_0, \cdots, q_n)$. From theorem 5.2.2, every 3-multisector $X_{(\mathbf{g})}$ is a twisted sector. Assume $X_{(\mathbf{g})} = \overline{O}_\tau = \mathbf{P}(Q_\tau)$, $\tau = (v_0, \cdots, v_{i-1})$ is a cone in the fan Ξ, $Q_\tau = (q_i, \cdots, q_n)$ and $gcd(q_i, \cdots, q_n) = d > 1$. So we have $g_1, g_2, g_3 \in \mathbf{Z}_d$ and $g_1 g_2 g_3 = 1$. $\mathbf{P}(Q_\tau) = \mathbf{P}^n(0, \cdots, 0, q_i, \cdots, q_n)$ is a hyperplane of $\mathbf{P}(Q)$. If we let $o(g_1) = l, o(g_2) = q, o(g_3) = r$, then $l + q + r \equiv 0 (mod\ d)$. From section 2.3., the fundamental group of orbifold $S^2(l, q, r)$ is:

$$\pi_1^{orb}(S^2) = \{\lambda_1, \lambda_2, \lambda_3 | \lambda_1^l = \lambda_2^q = \lambda_3^r = \lambda_1\lambda_2\lambda_3 = 1\}$$

Then $\Gamma = \pi_1^{orb}(S^2)$ is a fuchsian group. Let $\varphi : \Gamma \longrightarrow \mathbf{Z}_d$ be the homomorphism defined by $\lambda_i \longrightarrow g_i$ for $i = 1, 2, 3$. Assume $Im\varphi = G = K(\mathbf{g})$, there is a Riemann surface Σ such that $\Sigma / K(\mathbf{g}) = S^2(l, q, r)$. From [16], since the group $K(\mathbf{g})$ is also a cyclic group, we can describe the (l, q, r) as three cases:

(1) $(l, q, r) = (p, p, 0), p > 1$;
(2) $(l, q, r) = (3, 3, 3), (2, 3, 6), (2, 4, 4)$;
(3) (l, q, r) is not the cases of (1) and (2).

From (1) we can see that the Riemann surface $\Sigma = S^2$ and the cyclic group $\mathbf{Z}_p$ acts on S^2 with quotient $S^2(p, p, 0)$. In the case (2), the three groups of values are the only choices such that the Riemann surface Σ is the torus. In this case, $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} = 1$. In the case (3), we know that the universal covering space of $S^2(l, q, r)$ is $\mathbf{H}^2$(Poincare disk model),then $\Sigma = \mathbf{H}^2 / ker(\varphi)$. By Riemann-Hurwitz formula,

$$g(\Sigma) = \frac{|K(\mathbf{g})| + 2 - |K(\mathbf{g})|(\frac{1}{p} + \frac{1}{q} + \frac{1}{r})}{2} \tag{5.8}$$

In this case $\frac{1}{p} + \frac{1}{q} + \frac{1}{r} > 1$ and $g(\Sigma) \geq 2$.

Now we assume that the genus of Σ is g. Let $\gamma_1, \cdots, \gamma_{2g}$ is a canonical homological basis satisfying the following relations:

$$\begin{cases} \gamma_i \gamma_{g+i} = 1, \forall\ i = 1, \cdots, g; \\ \gamma_i \gamma_j = 0, \text{ otherwise.} \end{cases}$$

Then by Riemann bilinear relation [15], the period matrix is: $\left(\begin{smallmatrix} I_g \\ Z \end{smallmatrix}\right)$ such that:

$$\begin{cases} a.) & Z =^t Z; \\ b.) & ImZ \text{ is positively definitive.} \end{cases}$$

Suppose $\check{\gamma}_1, \cdots, \check{\gamma}_{2g}$ is the dual basis in $H^1(\Sigma)$, then there exists a basis $\omega_1, \cdots, \omega_g$ in $H^{0,1}(\Sigma)$ such that

$$(\omega_1, \cdots, \omega_g) = (\check{\gamma}_1, \cdots, \check{\gamma}_{2g}) \begin{pmatrix} I_g \\ Z \end{pmatrix}$$

The element λ_1 in Γ acts on $H^1(\Sigma)$ naturally. Let A^t be the representation matrix of $(\lambda_1)_*$ under the basis $\gamma_1, \cdots, \gamma_{2g}$, then λ_1^* acts on $H^1(\Sigma)$, the matrix representation is A. So:

$$\lambda_1^*(\omega_1, \cdots, \omega_g) = (\check{\gamma}_1, \cdots, \check{\gamma}_{2g}) \cdot A \cdot \begin{pmatrix} I_g \\ Z \end{pmatrix}.$$

We know that $A \cdot \binom{I_g}{Z}$ is a $2g \times g$ matrix. Assume the matrix which consists of the first g rows of the matrix $A \cdot \binom{I_g}{Z}$ is U. Since λ_1 is a holomorphic map, λ_1^* preserves the subspace $H^{1,0}(\Sigma)$. So

$$\lambda_1^*(\omega_1, \cdots, \omega_g) = (\omega_1, \cdots, \omega_g) \cdot U$$

We have

$$\lambda_1^*(\omega_1, \cdots, \omega_g) = (\check{\gamma}_1, \cdots, \check{\gamma}_{2g}) \cdot \begin{pmatrix} I_g \\ Z \end{pmatrix} \cdot U$$

Thus we can determine the matrix Z, and the matrix U can also be determined. We know that $H^{0,1}(\Sigma) = \overline{H^{1,0}(\Sigma)}$, so the basis of $\overline{H^{1,0}(\Sigma)}$ is $\overline{\omega}_1, \cdots, \overline{\omega}_g$, and

$$\lambda_1^*(\overline{\omega}_1, \cdots, \overline{\omega}_g) = (\overline{\omega}_1, \cdots, \overline{\omega}_g) \cdot \overline{U} \tag{5.9}$$

Let $U_j = \{[z]_Q \in \mathbf{P}^n_{q_0, \cdots, q_n} : z_j \neq 0\} \subset \mathbf{P}^n_{q_0, \cdots, q_n}$ for $j = 0, \cdots, n$. Then since $Q_\tau = (q_i, \cdots, q_n)$, we see that $X_{(\mathbf{g})} = \mathbf{P}(Q_\tau)$ can be covered by $X_{(\mathbf{g})} \cap U_i, \cdots, X_{(\mathbf{g})} \cap U_n$. From section 3.1, for $j \geq i$, we have a bijective map ϕ_j from U_j to $\mathbf{C}^n / \mu_{q_j}(Q_{q_j})$ given by

$$\phi_j([z]_Q) = \left(\frac{z_0}{(z_j)^{q_0/q_j}}, \cdots, \frac{\hat{z_j}}{z_j}, \cdots, \frac{z_n}{(z_j)^{q_n/q_j}} \right)_{q_j}$$

So we choose the coordinates of $\mathbf{C}^n = V_j$ by $\left(\frac{z_0}{(z_j)^{q_0/q_j}}, \cdots, 1, \cdots, \frac{z_n}{(z_j)^{q_n/q_j}} \right)$. If we let $x_0 = \frac{z_0}{(z_j)^{q_0/q_j}}, \cdots, x_j = 1, \cdots, x_n = \frac{z_n}{(z_j)^{q_n/q_j}}$, then let $p_j = [0, \cdots, 1, \cdots, 0]$ be the point in V_j, $(TV_j)_{p_j}$ has the basis $\frac{\partial}{\partial x_0}, \cdots, \frac{\partial}{\partial x_n}$, and the $\lambda_1^* = g_1$ acts on $(TV_j)_{p_j}$ in the natural way. We denote the diagonal representation matrix of λ_1^* by D. So on $(H^{0,1}(\Sigma) \otimes (TV_j)_{p_j})$ we have a basis:

$$\left\{ \frac{\partial}{\partial x_0} \otimes \overline{\omega}_1, \frac{\partial}{\partial x_1} \otimes \overline{\omega}_1, \cdots, \frac{\partial}{\partial x_n} \otimes \overline{\omega}_1, \frac{\partial}{\partial x_0} \otimes \overline{\omega}_2, \cdots, \frac{\partial}{\partial x_n} \otimes \overline{\omega}_n \right\}.$$

and $\lambda_1^* = g_1$ acts on $(H^{0,1}(\Sigma) \otimes (TV_j)_{p_j})$, which the matrix representation is $D \otimes \overline{U}$. Because g_1, g_2, g_3 generate the cyclic group $\mathbf{Z}_d$, and assume that the g_1 is a generator, so the matrix $D \otimes \overline{U}$ has eigenvalue 1 with multiplicity e, where

$$e = dim_{\mathbf{C}}(e(E_{(\mathbf{g})})) = dim_{\mathbf{C}}(X_{(\mathbf{g})}) - n + \Sigma_{j=1}^3 \iota_{(g_j)} \tag{5.10}$$

So locally the obstruction bundle is generated by the e eigenvectors $\xi_1, \cdots, \xi_e$.

Suppose $\xi = \alpha \otimes \omega \in \left((TV_j)_{p_j} \otimes H^{0,1}(\Sigma)\right)^{K(\mathbf{g})}$, then α is a linear combination of $\frac{\partial}{\partial x_0}, \cdots, \frac{\partial}{\partial x_{i-1}}$, otherwise ω will be a harmonic form on S^2 because ω is $K(\mathbf{g})$-invariant, so $\omega = 0$. And also from the action of $\lambda_1^* = g_1$ on $(TV_j)_{p_j} \otimes H^{0,1}(\Sigma)$ and $C(\mathbf{g})$ on the invariant subspace $((TV_j)_{p_j} \otimes H^{0,1}(\Sigma))^{\mathbf{Z}_d}$, we see that each eigenvector

ξ_t can be written as the linear combination of $\{\frac{\partial}{\partial x_t}\otimes\overline{\omega}_1,\cdots,\frac{\partial}{\partial x_t}\otimes\overline{\omega}_g\}$ for some t with $0\le t\le i-1$.

Now assume $k\ge i$, and $U_k\cap X_{(\mathbf{g})}$ is another open subset of $X_{(\mathbf{g})}$. Let $U_k=V_k/\mathbf{Z}_k$, from the above discussion, we can choose the coordinates of V_k as:

$$\left(y_0=\frac{z_0}{(z_k)^{q_0/q_k}},\cdots,y_k=1,\cdots,y_n=\frac{z_n}{(z_k)^{q_n/q_k}}\right).$$

So we have a basis on $(H^{0,1}(\Sigma)\otimes(TV_k)_{p_k})$:

$$\left\{\frac{\partial}{\partial y_0}\otimes\overline{\omega}_1,\frac{\partial}{\partial y_1}\otimes\overline{\omega}_1,\cdots,\frac{\partial}{\partial y_n}\otimes\overline{\omega}_1,\frac{\partial}{\partial y_0}\otimes\overline{\omega}_2,\cdots,\frac{\partial}{\partial y_n}\otimes\overline{\omega}_n\right\}.$$

Then we see that on the neighborhood $U_k\cap X_{(\mathbf{g})}$, the obstruction bundle is still generated by e eigenvectors $\xi_1^{'},\cdots,\xi_e^{'}$. And $\xi_l^{'}$ can be obtained from ξ_l by substituting $\frac{\partial}{\partial y_l}$ for $\frac{\partial}{\partial x_l}$. On the V_k, we have:

$$\begin{aligned}\frac{\partial}{\partial x_0}=\Sigma\frac{\partial y_j}{\partial x_0}\frac{\partial}{\partial y_j} &= \frac{\partial}{\partial x_0}(\frac{x_0}{(x_k)^{q_0/q_k}})\frac{\partial}{\partial y_0}\\ &= \frac{1}{(x_k)^{q_0/q_k}}\frac{\partial}{\partial y_0}\\ &= (y_j)^{q_0/q_j}\frac{\partial}{\partial y_0}\end{aligned}$$

For the same computation, we also have:

$$\begin{cases}\frac{\partial}{\partial x_1}=(y_j)^{q_1/q_j}\frac{\partial}{\partial y_1};\\ \cdots\cdots\\ \frac{\partial}{\partial x_{i-1}}=(y_j)^{q_{i-1}/q_j}\frac{\partial}{\partial y_{i-1}}.\end{cases}$$

Assume on the neighborhood $U_j\cap X_{(\mathbf{g})}$, the eigenvectors $\xi_1,\cdots,\xi_e$ are generated by the linear combination of the tuples $\{\frac{\partial}{\partial x_{t_1}}\otimes\overline{\omega}_1,\cdots,\frac{\partial}{\partial x_{t_1}}\otimes\overline{\omega}_g\},\cdots,\{\frac{\partial}{\partial x_{t_e}}\otimes\overline{\omega}_1,\cdots,\frac{\partial}{\partial x_{t_e}}\otimes\overline{\omega}_g\}$ respectively for $0\le t_1,\cdots,t_e\le i-1$. Then the transition function:

$$h_{kj}:(U_j\cap X_{(\mathbf{g})})\times\mathbf{C}^e\hookleftarrow(U_j\cap U_k\cap X_{(\mathbf{g})})\times\mathbf{C}^e\longrightarrow(U_j\cap U_k\cap X_{(\mathbf{g})})\times\mathbf{C}^e\hookrightarrow(U_k\cap X_{(\mathbf{g})})\times\mathbf{C}^e$$

can be written as:

$$h_{kj}(x;c_1,\cdots,c_e)=\left(x;y_j^{q_{t_1}/q_j}(x)\cdot c_1,\cdots,y_j^{q_{t_e}/q_j}(x)\cdot c_e\right).\tag{5.11}$$

So locally the transition matrix is:

$$\begin{pmatrix}y_j^{q_{t_1}/q_j}(x) & & \\ & \ddots & \\ & & y_j^{q_{t_e}/q_j}(x)\end{pmatrix}$$

Then the obstruction bundle $E_{(\mathbf{g})}$ can be splited as the whitney sum of line bundles, let $E_{(\mathbf{g})}=\oplus_{l=1}^{e}E_l$. Every line bundle E_l is generated by ξ_l on the neighborhood $U_j\cap X_{(\mathbf{g})}$. The group $\mathbf{Z}_d$ acts diagonally on the obstruction bundle $E_{(\mathbf{g})}$, so it acts on every line bundle E_l naturally. Assume the matrix representation of the action of the generator of $\mathbf{Z}_d$ on the obstruction bundle $E_{(\mathbf{g})}$ is:

$$\begin{pmatrix} e^{2\pi i\cdot\frac{m_1}{d}} & & \\ & \ddots & \\ & & e^{2\pi i\cdot\frac{m_e}{d}} \end{pmatrix}$$

here $0 \leq m_l < d, 1 \leq l \leq e$. Then we have the following facts. $e^{2\pi i\cdot\frac{m_l}{d}}$ is a d_l-root of 1 for $1 \leq l \leq e$, and it is clear that d_l is a divisor of d.

5.6. **Computation of the 3-Point Function.** In this section we use the localization technique ([3],[12]) to calculate the 3-point function defined in the orbifold cup product.

Let $X = \mathbf{P}(Q)$ be the weighted projective space of type $Q = (q_0, \cdots, q_n)$ and $X_{(\mathbf{g})}$ be a 3-multisector. Then $X_{(\mathbf{g})}$ is a twisted sector from theorem 5.2.2., assume $X_{(\mathbf{g})} = \overline{O}_\tau = \mathbf{P}(Q_\tau)$, where $Q_\tau = (0, \cdots, 0, q_i, \cdots, q_n)$,$\mathbf{g} = (g_1, g_2, g_3) \in T_3^0$ and $\tau = (v_0, \cdots, v_{i-1})$ is a cone of the fan $\Xi = \{v_0, \cdots, v_n\}$. The orbifold structure of $X_{(\mathbf{g})}$ can be described in Remark 4.3.6., From (2.4), the key calculation of the orbifold cup product is to calculate the 3-point function:

$$< \eta_1, \eta_2, \eta_3 >_{orb} = \int^{orb}_{X_{(\mathbf{g})}} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge e_A(E_{(\mathbf{g})}) \tag{5.12}$$

where $\eta_j \in H^*(X_{(g_j)}; \mathbf{Q})$, for $j = 1, 2, 3$.

Now we analyze the integral formula (5.12). In order to compute conveniently, we prove that we always can suppose $q_0 = 1$. If $q_0 \neq 1$, let $\widetilde{Q} = (1, q_0, \cdots, q_n)$, then $\mathbf{P}^n_{q_0,\cdots,q_n} \subset \mathbf{P}(\widetilde{Q}) = \mathbf{P}^{n+1}_{1,q_0,\cdots,q_n} = Y$ is a hypersurface which is obtained if by letting the first homogeneous coordinate of $\mathbf{P}(\widetilde{Q})$ be zero. From theorem 4.3.5 and theorem 5.2.2, it is easy to see that $\mathbf{P}(Q)$ and $\mathbf{P}(\widetilde{Q})$ have the same twisted sectors and 3-multisectors. Suppose that the matrix representations in $\mathbf{P}(\widetilde{Q})$ corresponding to g_1, g_2, g_3 in $\mathbf{P}(Q)$ are $\widetilde{g}_1, \widetilde{g}_2, \widetilde{g}_3$, then $Y_{(\widetilde{\mathbf{g}})} = Y_{(\widetilde{g}_1,\widetilde{g}_2,\widetilde{g}_3)} = X_{(\mathbf{g})}$. The cohomological classes $e_1^*\eta_1, e_2^*\eta_2, e_3^*\eta_3$ are invariant when they are taken as the cohomological classes of $Y_{(\widetilde{\mathbf{g}})}$. Suppose the homogeneous coordinates of $\mathbf{P}(\widetilde{Q})$ is $\mathbf{z} = [z, z_0, \cdots, z_n]$, let $\widetilde{U}_j = \{z_j \neq 0 | \mathbf{z} \in \mathbf{P}(\widetilde{Q})\}$, $(0 \leq j \leq n)$, $\widetilde{U} = \{z \neq 0 | \mathbf{z} \in \mathbf{P}(\widetilde{Q})\}$, then $Y_{(\widetilde{\mathbf{g}})}$ can be covered by $\bigsqcup_{j=i}^n \widetilde{U}_j \cap Y_{(\widetilde{\mathbf{g}})}$. For the local chart $\widetilde{U}_j$, let $\widetilde{U}_j = \widetilde{V}_j/\mathbf{Z}_{q_j}$ and choose the coordinates of $\widetilde{V}_j$ as $(x = \frac{z}{(z_j)^{1/q_j}}, x_0 = \frac{z_0}{(z_j)^{q_0/q_j}}, \cdots, x_n = \frac{z_n}{(z_j)^{q_n/q_j}})$, so we have a base of $(T\widetilde{V}_j)_{p_j}$: $(\frac{\partial}{\partial x}, \frac{\partial}{\partial x_0}, \cdots, \frac{\partial}{\partial x_n})$. Because the invariant subspace $((TV_j)_{p_j} \otimes H^{0,1}(\Sigma))^{\mathbf{Z}_d}$ is generated by $\xi_1, \cdots, \xi_e$, and we can see that the space $((TV_j)_{p_j} \otimes H^{0,1}(\Sigma))^{\mathbf{Z}_d} \subset ((T\widetilde{V}_j)_{\widetilde{p}_j} \otimes H^{0,1}(\Sigma))^{\mathbf{Z}_d}$. We construct a new obstruction bundle $E_{(\widetilde{\mathbf{g}})}$ over $Y_{(\widetilde{\mathbf{g}})}$ as follows. On the local chart $\widetilde{U}_j \cap Y_{(\widetilde{\mathbf{g}})}$, this bundle is given by $\widetilde{V}_j \cap H \times ((TV_j)_{p_j} \otimes H^{0,1}(\Sigma))^{\mathbf{Z}_d} \longrightarrow \widetilde{V}_j \cap H$, where $H = \{x = x_0 = \cdots = x_{i-1} = 0 | \mathbf{x} \in \widetilde{V}_j\}$ is a hypersurface of $\widetilde{V}_j$. It is easy to see that the transition function of this bundle is also given by (5.11), so it can also be splitted into the Whitney sum of line bundles. It is clear that $E_{(\widetilde{\mathbf{g}})} \cong E_{(\mathbf{g})}$, so we have

$$\int^{orb}_{X_{(\mathbf{g})}} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge e_A(E_{(\mathbf{g})}) = \int^{orb}_{Y_{(\widetilde{\mathbf{g}})}} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge e_{\widetilde{A}}(E_{(\widetilde{\mathbf{g}})}) \tag{5.13}$$

So in the following analysis, we assume that $q_0 = 1$, and we give a formula to compute the integration (5.12).

First from Remark 4.3.6, we know that if $d = gcd(q_i, \cdots, q_n) \neq 1$, $X_{(\mathbf{g})}$ is a nonreduced orbifold. From the discussion of section 5.5., let $E_{(\mathbf{g})} = \oplus_{l=1}^{e} E_l$, then for every line bundle E_l, using the same method of Park and Poddar [22], consider the associated orbifold principal bundle P_l of E_l such that $E_l = P_l \times_{S^1} \mathbf{C}$. We know that there is a global action of $\mathbf{Z}_{d_l}$ on each fibre $F = S^1$. The quotient $P_l/\mathbf{Z}_{d_l}$ is again an orbifold principal bundle over the orbifold $X_{(\mathbf{g})}$. Let $\pi_l : P_l \longrightarrow P_l/\mathbf{Z}_{d_l}$ be the quotient map, which extends to an orbifold bundle map. Choose an orbifold connection A_l that is the pullback $\pi_l^*(A_l^{'})$, where $A_l^{'}$ is an orbifold connection on the associated bundle $E_l^{'} = (P_l/\mathbf{Z}_{d_l}) \times_{S^1} \mathbf{C}$. The Lie algebra of F can be identified with $\mathbf{R}$, then the induced map on the lie algebra $(\pi_l)_* : \mathbf{R} \longrightarrow \mathbf{R}$ is just given by $a \longmapsto d_l a$.

Let Ω_l and $\Omega_l^{'}$ be the curvature 2-forms for A_l and $A_l^{'}$. By proposition 6.2 of [19], $(\pi_l)^*(\Omega_l^{'}) = d_l \Omega_l$. So

$$\int_{X_{(\mathbf{g})}}^{orb} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge e_A(E_{(\mathbf{g})}) = \int_{X_{(\mathbf{g})}}^{orb} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge \Pi_{l=1}^{e} e_{A_l}(E_l)$$

$$= \frac{1}{\Pi_l d_l} \int_{X_{(\mathbf{g})}}^{orb} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge \Pi_{l=1}^{e} e_{A_l^{'}}(E_l^{'}) \tag{5.14}$$

Since the action of $\mathbf{Z}_{d_l}$ in any uniformizing system of $E_l^{'}$ is trivial, $E_l^{'}$ induces an orbifold bundle $E_l^{''}$ over the reduced orbifold $X_{(\mathbf{g})}^{'}$ which has an induced connection $A_l^{''}$. The connections $A_l^{'}$ and $A_l^{''}$ may be represented by the same 1-form over V for $(V \times \mathbf{C}, G^{'}/\mathbf{Z}_{d_l}, \widetilde{\pi}_1^{''})$ of $E_l^{'}$ and $E_l^{''}$ respectively. By Chern-Weil theory, $\Omega_l^{'}$ and $\Omega_l^{''}$ can therefore be represented by the same 2-form on V. We know that $e_1^*(\eta_1), e_2^*(\eta_2)$ and $e_3^*(\eta_3)$ are invariant when taken as the cohomology classes of $X_{(\mathbf{g})}^{'}$. Since $K(\mathbf{g})$ acts on $X_{(\mathbf{g})}$ trivially, so from (2.1), we have:

$$\int_{X_{(\mathbf{g})}}^{orb} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge \Pi_{l=1}^{e} e_{A_l^{'}}(E_l^{'}) = \frac{1}{|\mathbf{Z}_d|} \int_{X_{(\mathbf{g})}^{'}}^{orb} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge \Pi_{l=1}^{e} e_{A_l^{''}}(E_l^{''}) \tag{5.15}$$

Next we mainly discuss the method to calculate the integration in (5.15). From above $X_{(\mathbf{g})} = \mathbf{P}(Q_\tau)$, $Q_\tau = (0, \cdots, 0, q_i, \cdots, q_n)$. So we have $X_{(\mathbf{g})}^{'} = \mathbf{P}(Q_\tau/d)$, $Q_\tau/d = (q_i/d, \cdots, q_n/d), d = gcd(q_i, \cdots, q_n)$. Let N_τ be the sublattice of N generated by $\tau = (v_0, \cdots, v_{i-1})$ and $N(\tau) = N/N_\tau$ be the quotient lattice. The fan of $X_{(\mathbf{g})}^{'}$ is given by the projection of Ξ to $N(\tau) \otimes \mathbf{R}$. The dual lattice of $N(\tau)$ is $M(\tau) = \tau^\perp \cap M$. The torus $T = spec(\mathbf{C}[M(\tau)]) = O_\tau$. The characters χ^m correspond to rational functions on $X_{(\mathbf{g})}^{'}$ when $m \in M(\tau)$. Then we can use the localization technique of [3], when reduced to orbifold, to calculate the integration (5.15).

We know $X_{(\mathbf{g})}^{'}$ is a toric variety, T acts on $X_{(\mathbf{g})}^{'}$ and this action has $n - i + 1$ fixed points p_j for $i \leq j \leq n$. We let $\{\rho_i\}$ be the basic characters of T action and $\{\lambda_i\}$ be the parameters of the lie algebra $t_{\mathbf{C}}$ of T corresponding to the above base $\{\rho_i\}$. Because $q_0 = 1$, it is clear that the matrix C_0 defined in proposition 3.2.2 is the unit matrix. We compute that the fan of the weighted projective space $\mathbf{P}(Q)$ is generated by $v_0, \cdots, v_n$, where $v_0 = (-q_1, \cdots, -q_n)$, $v_j = e_j$ for $1 \leq j \leq n$. Let

$\{m_1, \cdots, m_n\}$ be the standard basis of M, we calculate the base of $M(\tau)$ as:

$$\left\{\rho_1 = \frac{q_n}{d}m_i - \frac{q_i}{d}m_n, \cdots, \rho_r = \frac{q_n}{d}m_j - \frac{q_j}{d}m_n, \cdots, \rho_{n-i} = \frac{q_n}{d}m_{n-1} - \frac{q_{n-1}}{d}m_n\right\}$$

where $j = i + r - 1$. We first study the action of T on the normal bundle of p_j, i.e., the orbifold tangent space $(TX^{'}_{(\mathbf{g})})_{p_j}$.

Consider the fixed points $p_j (i \leq j \leq n-1)$. Denote the local coordinates on a uniformizing system of $X^{'}_{(\mathbf{g})}$ around p_j by: $[x_i, \cdots, 1, \cdots, x_n]$. Let $m^1 = a_1\rho_1 + \cdots + a_{n-i}\rho_{n-i}$, and $< m^1, v_i >= 1$, $< m^1, v_k >= 0$ for $k > i, k \neq j$. Then we have $a_r = -\frac{dq_i}{q_n q_j}, a_1 = \frac{d}{q_n}$, so $\chi^{m^1} = x_i$. Similarly, we compute $\chi^{m^t} = x_t$ for $t \neq r, n-i$, $m^t = \frac{d}{q_n}\rho_t - \frac{dq_{i+t-1}}{q_n q_j}\rho_r$. Using the same method, let $m^{n-i} = a_1\rho_1 + \cdots + a_{n-i}\rho_{n-i}$, and $< m^{n-i}, v_k >= 0$ for $k \geq i, k \neq j, n$, $< m^{n-i}, v_n >= 1$, we have $a_r = -\frac{d}{q_j}$, $m^{n-i} = -\frac{d}{q_j}\rho_r$, $\chi^{m^{n-i}} = x_n$. So the T-equivariant Euler class of normal bundle of $p_j (i \leq j \leq n)$ is given by

$$e_T\left(\nu_{p_j}\right) = \left(-\frac{d}{q_j}\lambda_r\right) \prod_{k \neq r} \frac{d}{q_n}\left(\lambda_k - \frac{q_{i+k-1}}{q_j}\lambda_r\right) \tag{5.16}$$

Now we consider the fixed point p_n, the local coordinates of the uniformizing system of $X^{'}_{(\mathbf{g})}$ is $[w_i, \cdots, w_{n-1}, 1]$. Using the same method, we obtain the T-equivariant Euler class of normal bundle of $p_n (i \leq j \leq n)$ is given by

$$e_T(\nu_{p_n}) = \prod_{k=1}^{n-i} \frac{d}{q_n}\lambda_k \tag{5.17}$$

Since $e_1^*(\eta_1), e_2^*(\eta_2)$ and $e_3^*(\eta_3)$ all belong to $H^*(X^{'}_{(\mathbf{g})}, \mathbf{Q})$. From the ordinary ring structure of weighted projective space in section 5.1, we only consider $\xi_1 \in H^2(X^{'}_{(\mathbf{g})}, \mathbf{Q})$, the generator of $H^2(X^{'}_{(\mathbf{g})}, \mathbf{Q}) = \mathbf{Q}$. Suppose $L \longrightarrow X^{'}_{(\mathbf{g})}$ be the canonical line bundle whose first chern-class is ξ_1. The corresponding Cartier divisior is $D = D_1 D_2 \cdots D_{i-1} D_i + \cdots + D_1 D_2 \cdots D_{i-1} D_n$, where $D_j = \{z_j = 0\} \subset \mathbf{P}(Q)$ is the basic divisor. Then from Oda [20], in the neighbor $U_j \cap X^{'}_{(\mathbf{g})}$, $(i \leq j \leq n-1)$, let $m = u_1\rho_1 + \cdots + u_{n-i}\rho_{n-i}$, and $< -m, v_i >= 1, \cdots$, $< -m, v_{j-1} >= 1$, $< -m, v_{j+1} >= 1, \cdots, < -m, v_n >= 1$, then we calculate $-m = \sum_{k \neq r} \frac{d}{q_n}\rho_k - \frac{d}{q_n}(\sum_{k \neq r} \frac{q_{k+i-1}}{q_j})\rho_r$, where $j = i + r - 1$. So the divisor D is given by the rational function χ^{-m} on $U_j \cap X^{'}_{(\mathbf{g})}$. Similarly, it is given by the rational function $\chi^{-m} = \chi^{\frac{d}{q_n}(\rho_1 + \cdots + \rho_{n-i})}$ on $U_n \cap X^{'}_{(\mathbf{g})}$. Hence the action of T on the corresponding line bundle of D at the fixed points p_j has weights

$$\text{On } p_j, (i \leq j \leq n-1): \sum_{k \neq r} \frac{d}{q_n}\lambda_k - \frac{d}{q_n}\left(\sum_{k \neq r} \frac{q_{k+i-1}}{q_j}\right)\lambda_r \tag{5.18}$$

$$\text{On } p_n: \frac{d}{q_n}\sum_k \lambda_k \tag{5.19}$$

On the other hand, we also can write $e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 = a(\xi_1)^s$, $a \in \mathbf{Q}$, s is an integer.

Now we analyze the Euler form $e(E^{''}_{(\mathbf{g})})$. From section 5.5., we compute the local generated vectors of the obstruction bundle $E_{(\mathbf{g})}$, and $E_{(\mathbf{g})} = \oplus_{l=1}^{e} E_l$. For each line bundle E_l, from (5.11), we have the transition function of E_l as

$$h_{jn}(x,c) = \left(x, x_n^{q_{t_l}/q_n}(x)\cdot c\right), \ (i \leq j \leq n-1)$$

Because the line bundle $E^{''}_l$ is the reduction of E_l under the $\mathbf{Z}_{d_l}$-invariant homomorphism, the transition function of the line bundle $E^{''}_l$

$$(U_n \cap X^{'}_{(\mathbf{g})}) \times c \supset (U_n \cap U_j \cap X^{'}_{(\mathbf{g})}) \times c \longrightarrow (U_n \cap U_j \cap X^{'}_{(\mathbf{g})}) \times c \subset (U_j \cap X^{'}_{(\mathbf{g})}) \times c$$

is given by

$$h_{jn}(x,c) = \left(x, (x_n^{q_{t_l}/q_n})^{d_l}(x)\cdot c\right), \ (i \leq j \leq n-1)$$

So we can define the action of T on $E^{''}_{(\mathbf{g})} = \bigoplus_{l=1}^{e} E^{''}_l$ as follows, on the line bundle $E^{''}_l$:

$$\begin{aligned}
&(1) \quad t(x,c) = (tx,c) = \left(tx, \chi^0(t)c\right), t \in T, (x,c) \in (U_n \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C};\\
&(2) \quad t(x,c) = \left(tx, (x_n^{q_{t_l}/q_n})^{d_l}(t)c\right) = \left(tx, (\chi^{-\frac{d}{q_j}\rho_r})^{\frac{q_{t_l}d_l}{q_n}}(t)c\right),\\
&\qquad t \in T, (x,c) \in (U_j \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C}.
\end{aligned}$$

where $(i \leq j \leq n-1), j = i+r-1$. Then the action of T on $E^{''}_l$ at the fixed points $p_n, p_j (i \leq j \leq n-1)$ has weights

$$0, \ -\frac{q_{t_l}d_l}{q_n} \cdot \frac{d}{q_j}\lambda_r. \tag{5.20}$$

So from the localization formula, when we consider the orbifold $X^{'}_{(\mathbf{g})}$, see Corollary 9.13 in [12], we have the integration

$$\int^{orb}_{X^{'}_{(\mathbf{g})}} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge \Pi_{l=1}^{e} e_{A^{''}_l}(E^{''}_l) = \frac{a\left(\frac{d}{q_n}\sum_{k=1}^{n-i}\lambda_k\right)^s \cdot 0}{a_n \cdot \prod_{k=1}^{n-i}\frac{d}{q_n}\lambda_k} +$$

$$\sum_{j=i}^{n-1} \frac{a\left[\sum_{k\neq r}\frac{d}{q_n}\lambda_k - \frac{d}{q_n}\left(\sum_{k\neq r}\frac{q_{k+i-1}}{q_j}\right)\lambda_r\right]^s \cdot \prod_{l=1}^{e}\left(-\frac{q_{t_l}d_l}{q_n}\cdot\frac{d}{q_j}\lambda_r\right)}{a_j \cdot \left(-\frac{d}{q_j}\lambda_r\right) \cdot \prod_{k\neq r}\frac{d}{q_n}\left(\lambda_k - \frac{q_{i+k-1}}{q_j}\lambda_r\right)} \tag{5.21}$$

Where $j = i+r-1$, a_j is the order of the local cyclic group of p_j in the orbifold $X^{'}_{(\mathbf{g})}$.

5.7. **Example.** In this example we use the methods of the above sections to calculate the 3-point functions. Let $Q = (1,2,2,3,3,3)$ and $\mathbf{P}(Q) = \mathbf{P}^5_{1,2,2,3,3,3}$ be the weighted projective space of type Q, then $q_0 = 1, q_1 = q_2 = 2, q_3 = q_4 = q_5 = 3$. From proposition 3.2.2., we have $C_0 = I_{5\times 5}$. So let $v_1 = e_1, v_2 = e_2, v_3 = e_3, v_4 = e_4, v_5 = e_5, v_0 = -\Sigma\frac{q_i}{q_0}v_i = (-2,-2,-3,-3,-3)$. The fan Ξ of $\mathbf{P}^5_{1,2,2,3,3,3}$ is generated by $\{v_0, v_1, v_2, v_3, v_4, v_5\}$. For $\sigma_5 = (v_0, v_1, v_2, v_3, v_4)$, we have $G_{\sigma_5} = N/N_{\sigma_5} =$

$\mathbf{Z}_3$. We write the matrix representation of the action of $\mathbf{Z}_3$ on $U_{\sigma_5'} = \mathbf{C}^5$ as follows.

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}; \begin{pmatrix} e^{2\pi i\cdot\frac{1}{3}} & 0 & 0 & 0 & 0 \\ 0 & e^{2\pi i\cdot\frac{2}{3}} & 0 & 0 & 0 \\ 0 & 0 & e^{2\pi i\cdot\frac{2}{3}} & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

$$\begin{pmatrix} e^{2\pi i\cdot\frac{2}{3}} & 0 & 0 & 0 & 0 \\ 0 & e^{2\pi i\cdot\frac{1}{3}} & 0 & 0 & 0 \\ 0 & 0 & e^{2\pi i\cdot\frac{1}{3}} & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

For $\sigma_4 = (v_0, v_1, v_2, v_3, v_5)$, and $\sigma_3 = (v_0, v_1, v_2, v_4, v_5)$, we have $G_{\sigma_4} = G_{\sigma_3} = \mathbf{Z}_3$. The actions of $\mathbf{Z}_3$ on $U_{\sigma_4'} = U_{\sigma_3'} = \mathbf{C}^5$ are the same as above.

For $\sigma_2 = (v_0, v_1, v_3, v_4, v_5)$. We write the matrix representation of the action of $\mathbf{Z}_2$ on $U_{\sigma_2'} = \mathbf{C}^5$ as follows.

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}; \begin{pmatrix} e^{2\pi i\cdot\frac{1}{2}} & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & e^{2\pi i\cdot\frac{1}{2}} & 0 & 0 \\ 0 & 0 & 0 & e^{2\pi i\cdot\frac{1}{2}} & 0 \\ 0 & 0 & 0 & 0 & e^{2\pi i\cdot\frac{1}{2}} \end{pmatrix}$$

For $\sigma_1 = (v_0, v_2, v_3, v_4, v_5)$, we have $G_{\sigma_1} = \mathbf{Z}_2$. The actions of $\mathbf{Z}_2$ on $U_{\sigma_1'} = \mathbf{C}^5$ is the same as above.

For $\sigma_0 = (v_1, v_2, v_3, v_4, v_5)$, $G_{\sigma_0} = 1$, and the action is trivial. If we let

$$g_1 = \begin{pmatrix} e^{2\pi i\cdot\frac{1}{3}} & 0 & 0 & 0 & 0 \\ 0 & e^{2\pi i\cdot\frac{2}{3}} & 0 & 0 & 0 \\ 0 & 0 & e^{2\pi i\cdot\frac{2}{3}} & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}; g_2 = \begin{pmatrix} e^{2\pi i\cdot\frac{1}{2}} & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & e^{2\pi i\cdot\frac{1}{2}} & 0 & 0 \\ 0 & 0 & 0 & e^{2\pi i\cdot\frac{1}{2}} & 0 \\ 0 & 0 & 0 & 0 & e^{2\pi i\cdot\frac{1}{2}} \end{pmatrix}$$

Then we have the twisted sectors: $X_{(g_1)} = X_{(g_1^2)} = \mathbf{P}(Q_\tau)$, where $Q_\tau = (0,0,0,3,3,3)$, $\tau = (v_0, v_1, v_2)$; $X_{(g_2)} = \mathbf{P}(Q_\delta)$, $Q_\delta = (0,2,2,0,0,0)$, $\delta = (v_0, v_3, v_4, v_5)$. The degree shifting numbers: $\iota_{(g_1)} = \frac{5}{3}$, $\iota_{(g_1^2)} = \frac{4}{3}$, $\iota_{(g_2)} = 2$. So the Chen-Ruan cohomology group of $\mathbf{P}^5_{1,2,2,3,3,3}$ is:

$$\begin{aligned} H^d_{orb}(\mathbf{P}^5_{1,2,2,3,3,3}; \mathbf{Q}) &= H^d(\mathbf{P}^5_{1,2,2,3,3,3}; \mathbf{Q}) \\ &\oplus H^{d-\frac{10}{3}}(\mathbf{P}(Q_\tau); \mathbf{Q}) \\ &\oplus H^{d-\frac{8}{3}}(\mathbf{P}(Q_\tau); \mathbf{Q}) \\ &\oplus H^{d-4}(\mathbf{P}(Q_\delta); \mathbf{Q}) \end{aligned}$$

All the 3-multisectors are: $X_{(g_1,g_1,g_1)} = X_{(g_1^2,g_1^2,g_1^2)} = \mathbf{P}(Q_\tau)$, $X_{(g_1,g_1^2,1)} = \mathbf{P}(Q_\tau)$, $X_{(g_2,g_2,1)} = \mathbf{P}(Q_\delta)$. In the 3-multisectors $X_{(g_1,g_1^2,1)}$ and $X_{(g_2,g_2,1)}$, from (5.2), the dimension of the obstruction bundle of these two 3-multisectors are all zero, so the integration (5.3) is the usual integration on orbifold. The orbifold cup product can be described easily.

For the 3-multisector $X_{(g_1^2,g_1^2,g_1^2)}$, the dimension of the obstruction bundle $E_{(\mathbf{g})}$ is 1. Let $X_{(\mathbf{g})} = X_{(g_1^2,g_1^2,g_1^2)}$, $\eta_j \in H^*(X_{(g_1^2)};\mathbf{Q}), (j = 1,2,3)$, then

$$< \eta_1, \eta_2, \eta_3 >_{orb}= \int_{X_{(\mathbf{g})}}^{orb} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge e_A(E_{(\mathbf{g})}) \tag{5.22}$$

Next we use the localization formula (5.21) to compute the 3-point function (5.22). First we describe the obstruction bundle $E_{(\mathbf{g})}$ over $X_{(\mathbf{g})}$. We see that g_1^2 generates the cyclic group $\mathbf{Z}_3$, so $K(\mathbf{g}) = \mathbf{Z}_3$. To describe the obstruction bundle, we consider the orbifold sphere $(S^2,(x_1,x_2,x_3),(3,3,3))$, the orbifold fundamental group is:

$$\pi_1^{orb}(S^2) = \left\{\lambda_1, \lambda_2, \lambda_3 | \lambda_i^3 = 1, \lambda_1\lambda_2\lambda_3 = 1\right\}.$$

Its orbifold universal cover is the Euclidean plane E^2([24]). We use the triangle group model $\triangle^*(p,q,r)$ for $\frac{1}{p}+\frac{1}{q}+\frac{1}{r} = 1$. Then from [24], the full triangle group $\triangle^*(p,q,r)$ is defined to be the triangle group of isometries of E^2 generated by the reflexions L,M and N in the three sides YZ,ZX and XY of $\triangle$. In each case, it is easy to see that the translates of $\triangle$ by $\triangle^*(3,3,3)$ title E^2. This tiling is shown in [24].

Now $\triangle^*(3,3,3)$ has a natural subgroup of index two-the orientation preserving subgroup which is denoted $\triangle(3,3,3)$. The product LM of two of the generating reflexions of $\triangle^*(3,3,3)$ is a rotation through $2\pi/3$ about Z and MN and NL are also rotations. Write $LM = \lambda_1$, $MN = \lambda_2$, $NL = \lambda_3$ so that $\lambda_1, \lambda_2, \lambda_3$ are rotations about Z, X, Y respectively, see Figure.1. Clearly $\lambda_1, \lambda_2, \lambda_3$ lie in $\triangle(3,3,3)$ and one can show that they generate it. So $\triangle(3,3,3) = \pi_1^{orb}(S^2)$, and $E^2/\pi_1^{orb}(S^2) = S^2(3,3,3)$. Let $P = \triangle \cup L\triangle$, then the $E^2/\pi_1^{orb}(S^2)$ is obtained from the fundamental region $\in= P$ by the following manner: XZ identified with LXZ, and XY identified with LXY.

Consider the homomorphism $\rho : \pi_1^{orb}(S^2) \longrightarrow K(\mathbf{g}) = \mathbf{Z}_3$ given by $\lambda_i \longmapsto g_1^2$. $ker(\rho)$ is a normal subgroup generated by commutators of λ_i and the element $\lambda_1\lambda_2^{-1}$. $ker(\rho)$ acts freely on E^2 with quotient being the hexigon shown in Figure.2.



Fig. 1



Fig. 2

Suppose $\alpha = \overrightarrow{ouw}$, $\beta = \overrightarrow{wvx}$. Then we obtain a symplectic basis $\{a,b\} = \{\alpha,\beta\}$ for $H_1(\Sigma;\mathbf{Z})$. Taking the Universal Coefficients Theorem dual, we obtain a canonical symplectic basis $\{\check{\alpha},\check{\beta}\}$ for $H^1(\Sigma;\mathbf{Z})$. From Riemann bilinear relation in [15], we know that there exists a complex basis ω for $H^{1,0}(\Sigma)$ such that its period matrix with respect to the a-class is the identity matrix and its period matrix with respect to the b-class is a symmetric complex matrix $R = (r)$ with ImR a positive matrix. Here we can write $\omega = \check{\alpha} + r\check{\beta}$.

The rotation λ_1 induces the following map on $\pi_1(\Sigma)$:

$$\begin{cases} \alpha \longmapsto \beta^{-1}; \\ \beta \longmapsto \alpha\beta^{-1}. \end{cases}$$

Hence the automorphism $(\lambda_1)_* : H_1(\Sigma) \longrightarrow H_1(\Sigma)$ can be expressed in the matrix:

$$\begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$$

with respect to the basis $\{\alpha, \beta\}$. The automorphism $\lambda_1^* : H^1(\Sigma) \longrightarrow H^1(\Sigma)$ can be expressed by the matrix:

$$A = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}$$

with respect to the dual basis $\{\check{\alpha}, \check{\beta}\}$, which is the transpose of the above matrix.

From the matrix A, we calculate that $\lambda_1^*\omega = -r\check{\alpha} + (-r+1)\check{\beta}$. Since λ_1 is a holomorphic map, λ_1^* preserves the subspace $H^{1,0}(\Sigma)$. Hence $\lambda_1^*\omega$ can be expressed as complex linear combination of ω, so we have $\lambda_1^*\omega = -r\omega$. We obtain $-r^2 = -r+1$, $r = \frac{1\pm\sqrt{3}i}{2}$. While $ImR > 0$, $r = \frac{1+\sqrt{3}i}{2}$. Because $H^{0,1}(\Sigma) = \overline{H^{1,0}(\Sigma)}$, $H^{0,1}(\Sigma)$ is generated by $\overline{\omega}$, $\overline{\omega} = \check{\alpha} + \bar{r}\check{\beta}$

Now let $\tau = (v_0, v_1, v_2)$, then $\overline{O}_\tau = X_{(\mathbf{g})}$. Suppose $U_i = \{z_i \neq 0\}$, then $X_{(\mathbf{g})} = (X_{(\mathbf{g})} \cap U_3) \cup (X_{(\mathbf{g})} \cap U_4) \cup (X_{(\mathbf{g})} \cap U_5)$. Let $U_3 = V_3/\mathbf{Z}_3$, $U_4 = V_4/\mathbf{Z}_3$, $U_5 = V_5/\mathbf{Z}_3$, from section 5.5., the coordinates of V_3,V_4 and V_5 are:

$$\begin{aligned} V_5 :\quad & \left\{x_0 = \tfrac{z_0}{(z_5)^{1/3}}, x_1 = \tfrac{z_1}{(z_5)^{2/3}}, x_2 = \tfrac{z_2}{(z_5)^{2/3}}, x_3 = \tfrac{z_3}{z_5}, x_4 = \tfrac{z_4}{z_5}, x_5 = 1\right\}; \\ V_4 :\quad & \left\{y_0 = \tfrac{z_0}{(z_4)^{1/3}}, y_1 = \tfrac{z_1}{(z_4)^{2/3}}, y_2 = \tfrac{z_2}{(z_4)^{2/3}}, y_3 = \tfrac{z_3}{z_4}, y_4 = 1, y_5 = \tfrac{z_5}{z_4}\right\}; \\ V_3 :\quad & \left\{w_0 = \tfrac{z_0}{(z_3)^{1/3}}, w_1 = \tfrac{z_1}{(z_3)^{2/3}}, w_2 = \tfrac{z_2}{(z_3)^{2/3}}, w_3 = 1, w_4 = \tfrac{z_4}{z_3}, w_5 = \tfrac{z_5}{z_3}\right\}. \end{aligned}$$

For the chart $(V_5, \mathbf{Z}_3, \pi_5)$, $TV_5|_{p_5}$ has framing $\{\frac{\partial}{\partial x_0}, \frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \frac{\partial}{\partial x_3}, \frac{\partial}{\partial x_4}\}$. So $TV_5|_{p_5} \otimes H^{0,1}(\Sigma)$ has framing

$$\left\{\frac{\partial}{\partial x_0} \otimes \overline{\omega}, \frac{\partial}{\partial x_1} \otimes \overline{\omega}, \frac{\partial}{\partial x_2} \otimes \overline{\omega}, \frac{\partial}{\partial x_3} \otimes \overline{\omega}, \frac{\partial}{\partial x_4} \otimes \overline{\omega}\right\}$$

Because $\lambda_1^*\overline{\omega} = -\overline{r\omega}$, and the action of λ_1^* on $TV_5|_{p_5}$ is expressed by the matrix g_1^2, so the representation matrix of the action of λ_1^* on $TV_5|_{p_5} \otimes H^{0,1}(\Sigma)$ is:

$$(*) \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{2} - \frac{\sqrt{3}}{2}i & 0 & 0 & 0 \\ 0 & 0 & -\frac{1}{2} - \frac{\sqrt{3}}{2}i & 0 & 0 \\ 0 & 0 & 0 & -\frac{1}{2} + \frac{\sqrt{3}}{2}i & 0 \\ 0 & 0 & 0 & 0 & -\frac{1}{2} + \frac{\sqrt{3}}{2}i \end{pmatrix}$$

So the matrix $(*)$ has eigenvalue 1 with multiplication one, and the corresponding eigenvector is $(1,0,0,0,0)$, the generator of $(TV_5|_{p_5} \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})}$ is $\xi_0 \otimes \overline{\omega} = \frac{\partial}{\partial x_0} \otimes \overline{\omega}$. Similarly, we compute the generator of $(TV_4|_{p_4} \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})}$ is $\xi_0^{'} \otimes \overline{\omega} = \frac{\partial}{\partial y_0} \otimes \overline{\omega}$ and the generator of $(TV_3|_{p_3} \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})}$ is $\xi_0^{''} \otimes \overline{\omega} = \frac{\partial}{\partial w_0} \otimes \overline{\omega}$.

Now we describe the local uniformizing charts for $E_{(\mathbf{g})}$:

- $\forall x \in X_{(\mathbf{g})}$, then $C(\mathbf{g}) = G_x = K(\mathbf{g}) = \mathbf{Z}_3$, and $(V_x^{\mathbf{g}} \times \mathbf{C}, K(\mathbf{g}), \widetilde{\pi})$ is a uniformizing system for $E_{(\mathbf{g})}$ where $K(\mathbf{g})$ acts on $V_x^{\mathbf{g}} \times \mathbf{C}$ by: $g_1^2(u,v) = (u, e^{2\pi i \cdot \frac{2}{3}} v)$.

The bundle $E_{(\mathbf{g})}$ is a line bundle, $dim_{\mathbf{C}} X_{(\mathbf{g})} = 2$, so the 3-point function (5.22) is nonzero only if there is some $\eta_i \in H^2(X_{(g_1)}; \mathbf{Q})$. Without loss of generality, assume $\eta_1 \in H^2(X_{(g_1^2)}; \mathbf{Q})$, $\eta_2 \in H^0(X_{(g_1^2)}; \mathbf{Q})$, $\eta_3 \in H^0(X_{(g_1^2)}; \mathbf{Q})$. In this case

$$< \eta_1, \eta_2, \eta_3 >_{orb} = \eta_2 \eta_3 \int^{orb}_{X_{(\mathbf{g})}} \eta_1 \wedge e_A(E_{(\mathbf{g})}) \tag{5.23}$$

From the first part of this section, we see that in this case, the orbifold principal S^1 bundle is $P_{(\mathbf{g})}$, and let $E^{'}_{(\mathbf{g})} = (P_{(\mathbf{g})}/K(\mathbf{g})) \times_{S^1} \mathbf{C}$ over $X_{(\mathbf{g})}$. $\pi_{K(\mathbf{g})} : P_{(\mathbf{g})} \longrightarrow P_{(\mathbf{g})}/K(\mathbf{g})$. Note that $\pi_{K(\mathbf{g})}$ on each fibre is given by $z \longmapsto z^3$. The lie algebra of $F = S^1$ can be identified with $\mathbf{R}$. Hence the induced map on the lie algebra $(\pi_{K(\mathbf{g})})_* : \mathbf{R} \longrightarrow \mathbf{R}$ is just multiplication by 3, so from (5.14)

$$\int^{orb}_{X_{(\mathbf{g})}} \eta_1 \wedge e_A(E_{(\mathbf{g})}) = \frac{1}{3} \int^{orb}_{X_{(\mathbf{g})}} \eta_1 \wedge e_{A'}(E^{'}_{(\mathbf{g})})$$

Where A and $A^{'}$ is the connections of $E_{(\mathbf{g})}$ and $E^{'}_{(\mathbf{g})}$ such that $\pi^*_{K(\mathbf{g})}(A^{'}) = A$. Then $E^{'}_{(\mathbf{g})}$ induces an orbifold bundle $E^{''}_{(\mathbf{g})}$ over the reduced orbifold $X^{'}_{(\mathbf{g})}$, from (5.15)

$$\int^{orb}_{X_{(\mathbf{g})}} \eta_1 \wedge e_{A'}(E^{'}_{(\mathbf{g})}) = \frac{1}{3} \int^{orb}_{X^{'}_{(\mathbf{g})}} \eta_1 \wedge e_{A''}(E^{''}_{(\mathbf{g})})$$

Where $A^{''}$ is the connection of $E^{''}_{(\mathbf{g})}$ induced from $E^{'}_{(\mathbf{g})}$. So we obtain

$$\int^{orb}_{X_{(\mathbf{g})}} \eta_1 \wedge e_A(E_{(\mathbf{g})}) = \frac{1}{9} \int^{orb}_{X^{'}_{(\mathbf{g})}} \eta_1 \wedge e_{A''}(E^{''}_{(\mathbf{g})}) \tag{5.24}$$

We now compute the integration $\int^{orb}_{X^{'}_{(\mathbf{g})}} \eta_1 \wedge e_{A''}(E^{''}_{(\mathbf{g})})$ in (5.24). The uniformizing system of $E^{''}_{(\mathbf{g})}$ over $X^{'}_{(\mathbf{g})}$ can be described as follows.

- $\forall x \in X^{'}_{(\mathbf{g})}$, then $C(\mathbf{g})/K(\mathbf{g}) = 1$ is the trivial group.

Now we use the localization technique to calculate the integration (5.24). Note that $X^{'}_{(\mathbf{g})} = O_\tau$, $\tau = (v_0, v_1, v_2)$, then $N(\tau) = N/N_\tau$, where N_τ is the sublattice generated by τ, and $M(\tau) = \tau^\perp \cap M$. The 2-torus associated to $X^{'}_{(\mathbf{g})}$ is $T = spec(\mathbf{C}[M(\tau)]) = O_\tau$. The characters χ^m correspond to rational functions on $X^{'}_{(\mathbf{g})}$ when $m \in M(\tau)$. Let $\{m_1, m_2, m_3, m_4, m_5\}$ be the standard basis of M. Then $\{\rho_1 = m_3 - m_5, \rho_2 = m_4 - m_5\}$ is a basis for $M(\tau)$. The T-action on $X^{'}_{(\mathbf{g})}$ has three fixed points p_3, p_4, p_5. First we study the action of T on the normal bundle of p_3, p_4 and p_5, i.e., the orbifold tangent space of p_3, p_4 and p_5.

From (5.16) and (5.17), of course we can compute using the same method as in section 5.6, we see that the T-equivariant Euler class of the normal bundle of p_5 is given by $e_T(\nu_{p_5}) = \lambda_1 \lambda_2$, and we also have: $e_T(\nu_{p_4}) = (\lambda_1 - \lambda_2)(-\lambda_2)$, $e_T(\nu_{p_3}) = (-\lambda_1 + \lambda_2)(-\lambda_1)$. In particular, we have $\chi^{-\rho_2} = y_5$ in the neighborhood V_4, and $\chi^{-\rho_1} = \omega_5$ in the neighborhood V_5.

The orbifold line bundle $E^{''}_{(\mathbf{g})}$ is trivialized by the generator $\frac{\partial}{\partial x_0} \otimes \overline{\omega}$ on $U_5 \cap X^{'}_{(\mathbf{g})}$, $\frac{\partial}{\partial y_0} \otimes \overline{\omega}$ on $U_4 \cap X^{'}_{(\mathbf{g})}$, and $\frac{\partial}{\partial w_0} \otimes \overline{\omega}$ on $U_3 \cap X^{'}_{(\mathbf{g})}$. On V_4: $\frac{\partial}{\partial x_0} = \sum \frac{\partial y_j}{\partial x_0} \frac{\partial}{\partial y_j} = \frac{\partial}{\partial x_0}(\frac{x_0}{x_4^{1/3}}) \frac{\partial}{\partial y_0} = y_5^{1/3} \frac{\partial}{\partial y_0}$; On V_3: $\frac{\partial}{\partial x_0} = \sum \frac{\partial w_j}{\partial x_0} \frac{\partial}{\partial w_j} = \frac{\partial}{\partial x_0}(\frac{x_0}{x_3^{1/3}}) \frac{\partial}{\partial w_1} = w_5^{1/3} \frac{\partial}{\partial w_0}$. So the transition function:

$h_{45} : (U_5 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C} \supset (U_4 \cap U_5 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C} \longrightarrow (U_4 \cap U_5 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C} \subset (U_4 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C}$ can be written as:

$$h_{45}(x,c) = \left(x, (y_5^{1/3})^3(x) \cdot c\right).$$

The transition function

$h_{35} : (U_5 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C} \supset (U_5 \cap U_3 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C} \longrightarrow (U_5 \cap U_3 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C} \subset (U_3 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C}$ is

$$h_{35}(x,c) = \left(x, (w_5^{1/3})^3(x) \cdot c\right)$$

So we can define the T action on $E^{''}_{(\mathbf{g})}$ by

(1) $t(x,c) = (tx, c) = (tx, \chi^0(t)c), t \in T, (x,c) \in (U_5 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C}$;
(2) $t(x,c) = (tx, y_5(t)c) = (tx, \chi^{-\rho_2}(t)c), t \in T, (x,c) \in (U_4 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C}$;
(3) $t(x,c) = (tx, w_5(t)c) = (tx, \chi^{-\rho_1}(t)c), t \in T, (x,c) \in (U_3 \cap X^{'}_{(\mathbf{g})}) \times \mathbf{C}$.

Then the action of T on $E^{''}_{(\mathbf{g})}$ at the fixed points p_5, p_4, p_3 has weights $0, (-\lambda_2), (-\lambda_1)$ respectively. we see that this can be calculated from (5.18) and (5.19).

Since $\eta_1 \in H^2(X_{(g_1^2)}; \mathbf{Q})$, so we can associate η_1 with a weil divisor $\{U_\sigma, \chi^{-m^\sigma}\}$. As before the transition functions $h_{\tau\sigma} : U_\sigma \times \mathbf{C} \supset U_{\sigma\cap\tau} \times \mathbf{C} \longrightarrow U_{\sigma\cap\tau} \times \mathbf{C} \subset U_\tau \times \mathbf{C}$ for the line bundle are given by $h_{\tau\sigma}(x,c) = (x, \chi^{(m_\sigma - m_\tau)}(x)c)$. Then one can define a T- action on the bundle that makes it a T-equivariant bundle ([20]) as follows:$t(x,c) = (tx, \chi^{-m_\sigma}(t)c)$ for $t \in T$ and $(x,c) \in U_\sigma \times \mathbf{C}$.

Suppose $D_i = \{z_i = 0\} (i = 0,1,2,3,4,5)$ are the divisors of X. We see that the divisor of $X^{'}_{(\mathbf{g})}$ can be expressed as the combination of $D_0D_1D_2D_3, D_0D_1D_2D_4$, $D_0D_1D_2D_5$. We consider the general η_1, and suppose it can be expressed as $D = a_3D_0D_1D_2D_3 + a_4D_0D_1D_2D_4 + a_5D_0D_1D_2D_5$. Then in $U_5 \cap X^{'}_{(\mathbf{g})}$, suppose $m = u_1\rho_1 + u_2\rho_2$, let $< -m, v_3 >= a_3, < -m, v_4 >= a_4$, then $u_1 = -a_3, u_2 = -a_4$, $\chi^{-m} = \chi^{a_3\rho_1 + a_4\rho_2}$. The divisor D is given by the rational function $\chi^{a_3\rho_1+a_4\rho_2}$ on $U_5 \cap X^{'}_{(\mathbf{g})}$. Similarly, it is given by the rational function $\chi^{-m} = \chi^{a_3\rho_1 - (a_5+a_3)\rho_2}$ on $U_4 \cap X^{'}_{(\mathbf{g})}$ and $\chi^{-(a_5+a_4)\rho_1 + a_4\rho_2}$ on $U_3 \cap X^{'}_{(\mathbf{g})}$. Hence the action of T on the corresponding line bundle of η_1 at the fixed points p_5, p_4 and p_3 has weights $a_3\lambda_1 + a_4\lambda_2, a_3\lambda_1 - (a_5+a_3)\lambda_2$ and $-(a_5+a_4)\lambda_1 + a_4\lambda_2$ respectively. So using the localization formula (5.21), we have:

$$\begin{aligned}
\int^{orb}_{X^{'}_{(\mathbf{g})}} \eta_1 \wedge e_{A''}(E^{''}_{(\mathbf{g})}) &= \frac{(a_3\lambda_1 + a_4\lambda_2)\cdot 0}{\lambda_1\lambda_2} + \frac{(a_3\lambda_1 - (a_5+a_3)\lambda_2)(-\lambda_2)}{(\lambda_1 - \lambda_2)(-\lambda_2)} \\
&+ \frac{(-(a_5+a_4)\lambda_1 + a_4\lambda_2)(-\lambda_1)}{(-\lambda_1 + \lambda_2)(-\lambda_1)} \\
&= (a_3 + a_4 + a_5)
\end{aligned}$$

So from (5.24), we have:

$$\int^{orb}_{X_{(\mathbf{g})}} \eta_1 \wedge e_A(E_{(\mathbf{g})}) = \frac{1}{9}(a_3 + a_4 + a_5).$$

From (5.23),

$$< \eta_1, \eta_2, \eta_3 >_{orb} = \frac{1}{9}(a_3 + a_4 + a_5)\eta_2\eta_3.$$

For example, if $D = D_0D_1D_2D_3$, then $a_3 = 1, a_4 = a_5 = 0$, so

$$< \eta_1, \eta_2, \eta_3 >_{orb} = \frac{1}{9}\eta_2\eta_3.$$

For the 3-multisector $X_{(g_1,g_1,g_1)} = \mathbf{P}(Q_\tau)$, the dimension of the obstruction bundle $E_{(\mathbf{g})}$ is 2. Let $X_{(\mathbf{g})} = X_{(g_1,g_1,g_1)}$, $\eta_j \in H^*(X_{(g_1)}; \mathbf{Q}), (j = 1, 2, 3)$, then

$$< \eta_1, \eta_2, \eta_3 >_{orb} = \int^{orb}_{X_{(\mathbf{g})}} e_1^*\eta_1 \wedge e_2^*\eta_2 \wedge e_3^*\eta_3 \wedge e_A(E_{(\mathbf{g})}) \tag{5.25}$$

Next we use the localization technique to calculate the 3-point function (5.25). First we describe the obstruction bundle $E_{(\mathbf{g})}$ over $X_{(\mathbf{g})}$. We know that g_1 generate the group $\mathbf{Z}_3$, so $K(\mathbf{g}) = \mathbf{Z}_3$. We consider the orbifold sphere $(S^2, (x_1, x_2, x_3), (3, 3, 3))$, let

$$\rho : \pi_1^{orb}(S^2) \longrightarrow K(\mathbf{g}) = \mathbf{Z}_3$$

is the surjective homomorphism defined by $\lambda_i \longmapsto g_1$. Then we have a Riemann surface $\Sigma = E^2/ker(\rho)$. Using the same method above, we have a basis $\overline{\omega}$ of $H^{0,1}(\Sigma)$, and

$$\lambda_1^*(\overline{\omega}) = -\overline{r\omega}$$

Let $\tau = (v_0, v_1, v_2)$, then we know that $\overline{O}_\tau = X_{(\mathbf{g})}$. Assume $U_i = \{z_i \neq 0\}$, then $X_{(\mathbf{g})} = (X_{(\mathbf{g})} \cap U_3) \cup (X_{(\mathbf{g})} \cap U_4) \cup (X_{(\mathbf{g})} \cap U_5)$. Write $U_3 = V_3/\mathbf{Z}_3$, $U_4 = V_4/\mathbf{Z}_3$, $U_5 = V_5/\mathbf{Z}_3$, then we can choose the coordinates of the open set V_3, V_4 and V_5 the same as before.

For the coordinate neighborhood $(V_5, \mathbf{Z}_3, \pi_5)$, $TV_5|_{p_5}$ has a framing: $\{\frac{\partial}{\partial x_0}, \frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \frac{\partial}{\partial x_3}, \frac{\partial}{\partial x_4}\}$, so $TV_5|_{p_5} \otimes H^{0,1}(\Sigma)$ has a framing

$$\left\{\frac{\partial}{\partial x_0} \otimes \overline{\omega}, \frac{\partial}{\partial x_1} \otimes \overline{\omega}, \frac{\partial}{\partial x_2} \otimes \overline{\omega}, \frac{\partial}{\partial x_3} \otimes \overline{\omega}, \frac{\partial}{\partial x_4} \otimes \overline{\omega}\right\}$$

Because $\lambda_1^*\overline{\omega} = -\overline{r\omega}$, and the action λ_1^* on $TV_5|_{p_5}$ can be represented by the matrix g_1, so the representation matrix of the action of λ_1^* on $TV_5|_{p_5} \otimes H^{0,1}(\Sigma)$ is:

$$(**) \begin{pmatrix} -\frac{1}{2} - \frac{\sqrt{3}}{2}i & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -\frac{1}{2} + \frac{\sqrt{3}}{2}i & 0 \\ 0 & 0 & 0 & 0 & -\frac{1}{2} + \frac{\sqrt{3}}{2}i \end{pmatrix}$$

We see that the matrix $(**)$ has eigenvalue 1 with multiplication 2, the corresponding eigenvectors are: $(0, 1, 0, 0, 0)$ and $(0, 0, 1, 0, 0)$, so the invariant subspace $(TV_5|_{p_5} \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})}$ has generators: $\xi_1 \otimes \overline{\omega} = \frac{\partial}{\partial x_1} \otimes \overline{\omega}$ and $\xi_2 \otimes \overline{\omega} = \frac{\partial}{\partial x_2} \otimes \overline{\omega}$. Similarly, $(TV_4|_{p_4} \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})}$ has generators: $\xi_1^{'} \otimes \overline{\omega} = \frac{\partial}{\partial y_1} \otimes \overline{\omega}$ and $\xi_2^{'} \otimes \overline{\omega} = \frac{\partial}{\partial y_2} \otimes \overline{\omega}$; $(TV_3|_{p_3} \otimes H^{0,1}(\Sigma))^{K(\mathbf{g})}$ has generators: $\xi_1^{''} \otimes \overline{\omega} = \frac{\partial}{\partial w_1} \otimes \overline{\omega}$ and $\xi_2^{''} \otimes \overline{\omega} = \frac{\partial}{\partial w_2} \otimes \overline{\omega}$.

We describe the uniformizing system of $E_{(\mathbf{g})}$ as follows.

- $\forall x \in X_{(\mathbf{g})}$, then $C(\mathbf{g}) = G_x = K(\mathbf{g}) = \mathbf{Z}_3$, and if $(V_x^{\mathbf{g}} \times \mathbf{C}^2, K(\mathbf{g}), \widetilde{\pi})$ is a uniformizing system of the bundle $E_{(\mathbf{g})}$, then $K(\mathbf{g})$ acts on $V_x^{\mathbf{g}} \times \mathbf{C}^2$ through: $g_1(u, v_1, v_2) = (u, e^{2\pi i \cdot \frac{2}{3}}v_1, e^{2\pi i \cdot \frac{2}{3}}v_2)$.

The obstruction bundle $E_{(\mathbf{g})}$ is a plane bundle. And $dim_{\mathbf{C}} X_{(\mathbf{g})} = 2$, so the 3-point function (5.25) is nonzero only if $\eta_j \in H^0(X_{(g_1)}; \mathbf{Q}), j = 1, 2, 3$. In this case

$$< \eta_1, \eta_2, \eta_3 >_{orb} = \eta_1\eta_2\eta_3 \int_{X_{(\mathbf{g})}}^{orb} e_A(E_{(\mathbf{g})}) \tag{5.26}$$

From the section 5.5., the obstruction bundle $E_{(\mathbf{g})}$ is the whitney sum of two orbifold line bundles, let $E_{(\mathbf{g})} = E_1 \oplus E_2$. E_1 is generated by $\xi_1 \otimes \overline{\omega}$ on the negiborhood $U_5 \cap X_{(\mathbf{g})}$; and E_2 is generated by $\xi_2 \otimes \overline{\omega}$ on $U_5 \cap X_{(\mathbf{g})}$. So from the first part of this section, consider the orbifold principal S^1-bundle P_l of $E_l (l = 1, 2)$. From the section 5.5, we can see that $\mathbf{Z}_{d_l} = K(\mathbf{g}) = \mathbf{Z}_3$, so let $E_l^{'} = (P_l/K(\mathbf{g})) \times_{S^1} \mathbf{C}$ is the orbifold bundle over $X_{(\mathbf{g})}$. $\pi_{K(\mathbf{g})} : P_l \longrightarrow P_l/K(\mathbf{g})$ is the projective map. Note that on every fibre, $\pi_{K(\mathbf{g})}$ is given by: $z \longmapsto z^3$. The lie algebra of $F = S^1$ is $\mathbf{R}$. So the induced map on the lie algebra is: $(\pi_{K(\mathbf{g})})_* : \mathbf{R} \longrightarrow \mathbf{R}, a \longmapsto 3a$. From (5.14),

$$\int_{X_{(\mathbf{g})}}^{orb} e_A(E_{(\mathbf{g})}) = \frac{1}{9} \int_{X_{(\mathbf{g})}}^{orb} \Pi_{l=1}^2 e_{A_l^{'}}(E_l^{'})$$

where A_l and $A_l^{'}$ are the connections on the bundles E_l and $E_l^{'}$ such that $\pi_{K(\mathbf{g})}^*(A_l^{'}) = A_l$. In this moment the group $K(\mathbf{g})$ acts on the bundle $E_l^{'}$ trivially, so $E_l^{'}$ induced an orbifold bundle $E_l^{''}$ over the reduced orbifold $X_{(\mathbf{g})}^{'}$. From (5.15),

$$\int_{X_{(\mathbf{g})}}^{orb} \Pi_{l=1}^2 e_{A_l^{'}}(E_l^{'}) = \frac{1}{3} \int_{X_{(\mathbf{g})}^{'}}^{orb} \Pi_{l=1}^2 e_{A_l^{''}}(E_l^{''})$$

where $A_l^{''}$ is the connection on the bundle $E_l^{''}$ induced from the bundle $E_l^{'}$. Thus

$$\int_{X_{(\mathbf{g})}}^{orb} e_A(E_{(\mathbf{g})}) = \frac{1}{27} \int_{X_{(\mathbf{g})}^{'}}^{orb} \Pi_{l=1}^2 e_{A_l^{''}}(E_l^{''}) \tag{5.27}$$

We now calculate the integration $\int_{X_{(\mathbf{g})}^{'}}^{orb} \Pi_{l=1}^2 e_{A_l^{''}}(E_l^{''})$. The unformizing system of the bundle $E_l^{''}$ over the reduced orbifold $X_{(\mathbf{g})}^{'}$ can be described as:

- For $\forall x \in X_{(\mathbf{g})}^{'}$, then $C(\mathbf{g})/K(\mathbf{g}) = 1$ is the trivial group, and the action is trivial.

Now we use the localization technique to compute the integration (5.27). We know that $X_{(\mathbf{g})}^{'} = \overline{O}_\tau$ is a toric variety, $\tau = (v_0, v_1, v_2)$. The three fixed points by the T-action on $X_{(\mathbf{g})}^{'}$ are p_3, p_4, p_5. We already computed the T-equivariant Euler class of the point p_5 at the normal bundle is:

$$e_T(\nu_{p_5}) = \lambda_1\lambda_2$$

Similarly, the T-equivariant Euler classes of the points p_4 and p_3 at the normal bundles are:

$$e_T(\nu_{p_4}) = (\lambda_1 - \lambda_2)(-\lambda_2), e_T(\nu_{p_3}) = (-\lambda_1 + \lambda_2)(-\lambda_1).$$

Orbifold line bundle $E_l^{''}, (l = 1, 2)$ is trivialized by $\frac{\partial}{\partial x_l} \otimes \overline{\omega}, (l = 1, 2)$ on $U_5 \cap X_{(\mathbf{g})}^{'}$, $\frac{\partial}{\partial y_l} \otimes \overline{\omega}, (l = 1, 2)$ on $U_4 \cap X_{(\mathbf{g})}^{'}$, and $\frac{\partial}{\partial w_l} \otimes \overline{\omega}, (l = 1, 2)$ on $U_3 \cap X_{(\mathbf{g})}^{'}$. On the neighborhood V_4, we have:

$$\begin{aligned}\xi_1 = \frac{\partial}{\partial x_1} = \sum \frac{\partial y_j}{\partial x_1}\frac{\partial}{\partial y_j} &= \frac{\partial}{\partial x_1}\Big(\frac{x_1}{x_4^{2/3}}\Big)\frac{\partial}{\partial y_1} \\ &= y_5^{2/3}\frac{\partial}{\partial y_1}\end{aligned}$$

Similarly, we can compute $\xi_2 = \frac{\partial}{\partial x_2} = y_5^{2/3}\frac{\partial}{\partial y_2}$.

On the neighborhood V_3, we have:

$$\begin{aligned}\xi_1 = \frac{\partial}{\partial x_1} = \sum \frac{\partial w_j}{\partial x_1}\frac{\partial}{\partial w_j} &= \frac{\partial}{\partial x_1}\Big(\frac{x_1}{x_3^{2/3}}\Big)\frac{\partial}{\partial w_1} \\ &= w_5^{2/3}\frac{\partial}{\partial w_1}\end{aligned}$$

Similarly, we have $\xi_2 = \frac{\partial}{\partial x_2} = w_5^{2/3}\frac{\partial}{\partial w_2}$. So the transition function of the orbifold line bundle $E_l^{''}, (l = 1, 2)$

$h_{45} : (U_5 \cap X'_{(\mathbf{g})}) \times \mathbf{C} \supset (U_4 \cap U_5 \cap X'_{(\mathbf{g})}) \times \mathbf{C} \longrightarrow (U_4 \cap U_5 \cap X'_{(\mathbf{g})}) \times \mathbf{C} \subset (U_4 \cap X'_{(\mathbf{g})}) \times \mathbf{C}$

is:

$$h_{45}(x, c) = \Big(x, (y_5^{2/3})^3(x) \cdot c\Big)$$

The transition function

$h_{35} : (U_5 \cap X'_{(\mathbf{g})}) \times \mathbf{C} \supset (U_5 \cap U_3 \cap X'_{(\mathbf{g})}) \times \mathbf{C} \longrightarrow (U_5 \cap U_3 \cap X'_{(\mathbf{g})}) \times \mathbf{C} \subset (U_3 \cap X'_{(\mathbf{g})}) \times \mathbf{C}$

is:

$$h_{35}(x, c) = \Big(x, (w_5^{2/3})^3(x) \cdot c\Big)$$

So we can define the action of T on the bundle $E_l^{''}$ as

(1) $t(x, c) = (tx, c) = (tx, \chi^0(t)c), t \in T, (x, c) \in (U_5 \cap X'_{(\mathbf{g})}) \times \mathbf{C}$;
(2) $t(x, c) = (tx, y_5^2(t)c) = (tx, \chi^{-2\rho_2}(t)c), t \in T, (x, c) \in (U_4 \cap X'_{(\mathbf{g})}) \times \mathbf{C}$;
(3) $t(x, c) = (tx, w_5^2(t)c) = (tx, \chi^{-2\rho_1}(t)c), t \in T, (x, c) \in (U_3 \cap X'_{(\mathbf{g})}) \times \mathbf{C}$.

Then the action of T on $E_l^{''}$ at the fixed points p_5, p_4, p_3 has weights $0, (-2\lambda_2), (-2\lambda_1)$ respectively.

So from (5.21), we have:

$$\begin{aligned}\int^{orb}_{X'_{(\mathbf{g})}} \Pi^2_{t=1} e_{A''_t}(E''_t) &= \frac{0}{\lambda_1\lambda_2} + \frac{(-2\lambda_2)^2}{(\lambda_1 - \lambda_2)(-\lambda_2)} + \frac{(-2\lambda_1)^2}{(-\lambda_1 + \lambda_2)(-\lambda_1)} \\ &= 4\end{aligned}$$

From (5.27),

$$\int^{orb}_{X_{(\mathbf{g})}} e_A(E_{(\mathbf{g})}) = \frac{4}{27}.$$

And by (5.26),

$$< \eta_1, \eta_2, \eta_3 >_{orb} = \frac{4}{27}\eta_1\eta_2\eta_3.$$

Acknowledgments. We thank Professor Banghe Li, Yongbin Ruan , jianzhong Pan, David Cox, and Mainak Poddar for very helpful encouragements. We especially thank Professor Yongbin Ruan for explaining to me to consider this interesting problem.

.

References

[1] Abd' Allah Al Amrami, *Cohomological study of weighted projective space*, In Algebraic Geometry, Lecture Notes in pure and applied mathematics, 193, Edited by Sinan Sertoz.

[2] W.Baily, Jr, The decomposition theorem for V-manifold, *Amer. J. Math* 78 (1956). (1984), 1-28.

[3] M.Atiyah and R.Bott, The moment map and equivariant cohomology, *Topology* 23 (1984), 1-28.

[4] Chen Hao, Chen-Ruan orbifold cohomology of moduli space $\mathcal{M}^{0,n}/S_n$ and complex dynamic systems, Ph.D thesis.

[5] H.Conrads, Weighted projective spaces and reflexive simplices, *Manuscripta. Mathematica*,Vol 107, No.2, 2002, 215-227.

[6] W.Chen and Y.Ruan, A new cohomology theory for orbifolds, math AG/0004129.

[7] W.Chen and Y.Ruan, Orbifold Gromov-Witten theory,*Contem. Math.* 310, math.AG/0103156.

[8] D.Cox, The homogeneous coordinate ring of a toric variety, *J. Algebraic Geometry* 4 (1995), 17-50, alg-geom/9210008.

[9] D.Cox, What is atoric variety? preprint.

[10] L.Dixon, J.Harvey, C.Vafa, E.Witten, Strings on orbifolds I, *Nucl. phys.* B 261, 678-686 (1985).

[11] L.Dixon, J.Harvey, C.Vafa, E.Witten, Strings on orbifolds II, *Nucl. phys.* B 274, 285-314 (1986).

[12] David A.Cox, Sheldon Katz, *Mirror Symmetry and Algebraic Geometry*, Mathematical Surveys and Monographs, Volume 68.

[13] W.Fulton, *Introduction to Toric Varieyies*, Annals of Mathematics Studies, 131, The William H. Roever Lectures in Geometry, Princeton Univ. Press, Princeton, NJ. 1993.

[14] Leonor Godinho, Blowing up Symplectic Orbifolds, *Annals of Global Analysis and Geometry*, 20, No.2 (2001), 117-162.

[15] P.A.Griffths, *Introduction to Algebraic Curves*, Translated from the Chinese by Kuniko Weltin, Translations of Mathematical Monographs, 76, Amer. Math. Soc., Providence, RI, 1989.

[16] W.J.Harvey, Cyclic group of automorphisms of a compact Riemann Surface, *Quart. J. Math.* 17 (1966), 86-97.

[17] Kawaski,T., Cohomology of twisted projective spaces and lens complexes, *Math. Ann.* 206 (1973) 243-248.

[18] Kawaski,T., The signature theorem for V-manifolds, *Topology* 17 (1978), 75-83.

[19] S.Kobayashi and K.Nomizu, *Foundations of Differential Geometry*, Vol.I, Interscience Publishers, a division of John Wiley and sons, New York, 1963.

[20] T.Oda, *Covex Bodies and Algebraic Geometry*, Springer-Verlag, Berlin-Heidelberg- New York, 1988.

[21] Mainak Poddar, Orbifold Hodge numbers of Calabi-Yau hypersurfaces, *Pacific J. Math.* Vol. 208, No.1, (2003), 151-167.

[22] B.Doug Park and Mainak Poddar, The Chen-Ruan Cohomology ring of Mirror Quintic, preprint.

[23] Y. Ruan, Cohomology ring of crepant resolution of orbifolds, math.AG/0108195.

[24] P.Scott, The geometries of 3-manifolds, *Bull. London. Math. Soc.* 15 (1983) No.5, 401-487.

[25] E.Zaslow, Topological orbifold models and quantum cohomology rings, *Comm. Math. Phys.* 156 (1993), No.2, 301-331.

Institute of Systems Science, Academy of Mathematics and System Sciences, Chinese Academy of Sciences, Beijing, 100080, P.R.China
E-mail address: `jiangyf@mail.amss.ac.cn`